

Product Data Sheet

MED12L siRNA (Human)

Reactivity	Applications		
tic H	RNAi		
Description siRNA to inhibit MED12L expression using RNA interference			
MED12L siRNA (Hur	man) is a target-specific 19-23 nt siRNA oligo duplexes designed		
to knock down gene	e expression.		
Lyophilized powder			
MED12L			
KIAA1635; TNRC111	; TRALP; TRALPUSH; Mediator of RNA polymerase II		
transcription subun	it 12-like protein; Mediator complex subunit 12-like protein;		
Thyroid hormone re	eceptor-associated-like protein; Trinucleotide repeat-containing		
gene 11 protein-like			
Entrez Gene 116931 (Human)			
SwissProt Q86YW9 (Human)			
> 97%			
Oligonucleotide syn	thesis is monitored base by base through trityl analysis to ensure		
appropriate couplin	g efficiency. The oligo is subsequently purified by affinity-solid		
phase extraction. Th	ne annealed RNA duplex is further analyzed by mass		
spectrometry to ver	rify the exact composition of the duplex. Each lot is compared to		
the previous lot by	mass spectrometry to ensure maximum lot-to-lot consistency.		
We offers pre-desig	ned sets of 3 different target-specific siRNA oligo duplexes of		
human MED12L ger	ne. Each vial contains 5 nmol of lyophilized siRNA. The duplexes		
can be transfected i	ndividually or pooled together to achieve knockdown of the		
target gene, which i	is most commonly assessed by qPCR or western blot.		
	tic H siRNA to inhibit ME MED12L siRNA (Hur to knock down gene Lyophilized powder MED12L KIAA1635; TNRC11H transcription subun Thyroid hormone re gene 11 protein-like 116931 (Human) Q86YW9 (Human) > 97% Oligonucleotide syn appropriate couplin phase extraction. Th spectrometry to ven the previous lot by We offers pre-desig human MED12L gen can be transfected i		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

Component	15 nmol	30 nmol
MED12L siRNA (Human) - A	5 nmol x 1	5 nmol x 2
MED12L siRNA (Human) - B	5 nmol x 1	5 nmol x 2
MED12L siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μΙ
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
12-well 1 ml		100 nM	5 µl	2 μl
	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
6-well	2 ml	100 nM	10 µl	5 μl
	2 ml	50 nM	5 μl	5 μΙ

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com