

# **Product Data Sheet**

## GALNT15 siRNA (Human)

Catalog #	Source	Reactivity	Applications		
CRJ4565	Synthetic	Н	RNAi		
Description	siRNA	to inhibit GALNT15	expression using RNA interference		
Specificity	GALN	T15 siRNA (Human) i	s a target-specific 19-23 nt siRNA oligo duplexes designed		
	to kno	ock down gene expre	ession.		
Form	Lyoph	ilized powder			
Gene Symbol	GALN	GALNT15			
Alternative Na	ames GALN	GALNTL2; Polypeptide N-acetylgalactosaminyltransferase 15; Polypeptide GalNAc			
	transf	ferase-like protein 2;	GalNAc-T-like protein 2; pp-GaNTase-like protein 2;		
	Polyp	eptide N-acetylgalac	tosaminyltransferase-like protein 2; Protein-UDP		
	acety	lgalactosaminyltra			
Entrez Gene	11724	117248 (Human)			
SwissProt	Q8N3	Q8N3T1 (Human)			
Purity	> 97%	, D			
Quality Contro	ol Oligoi	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure			
	appro	priate coupling effic	ency. The oligo is subsequently purified by affinity-solid		
	phase	e extraction. The ann	ealed RNA duplex is further analyzed by mass		
	spect	rometry to verify the	exact composition of the duplex. Each lot is compared to		
	the p	revious lot by mass s	pectrometry to ensure maximum lot-to-lot consistency.		
Components	We of	ffers pre-designed se	ts of 3 different target-specific siRNA oligo duplexes of		
	huma	n GALNT15 gene. Ea	ch vial contains 5 nmol of lyophilized siRNA. The duplexes		
	can b	e transfected individ	ually or pooled together to achieve knockdown of the		
	target	t gene, which is most	commonly assessed by qPCR or western blot.		
		And the second s	abiata abaya internet IC. Jacobs and fluores and a CC. Floures at a state state of C		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

Component	15 nmol	30 nmol
GALNT15 siRNA (Human) - A	5 nmol x 1	5 nmol x 2
GALNT15 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
GALNT15 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 µl	2 μΙ
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
6-well	2 ml	100 nM	10 µl	5 µl
6-Well	2 ml	50 nM	5 μl	5 μΙ

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com