

Product Data Sheet

SLC34A3 siRNA (Human)

| Catalog # | Source | Reactivity | Applications | | | |
|----------------|-----------|---|--|--|--|--|
| CRJ5352 | Synthetic | Н | RNAi | | | |
| Description | siRNA | A to inhibit SLC34A3 e | xpression using RNA interference | | | |
| Specificity | SLC34 | 4A3 siRNA (Human) is | a target-specific 19-23 nt siRNA oligo duplexes designed | | | |
| | to kn | ock down gene expre | ssion. | | | |
| Form | Lyopł | nilized powder | | | | |
| Gene Symbol | SLC34 | SLC34A3 | | | | |
| Alternative Na | ames NPT2 | NPT2C; NPTIIC; Sodium-dependent phosphate transport protein 2C; | | | | |
| | Sodiu | ım-phosphate transpo | ort protein 2C; Na(+)-dependent phosphate cotransporter | | | |
| | 2C; S | odium/inorganic pho | sphate cotransporter IIC; Sodium/phosphate | | | |
| | cotra | nsporter 2C; Na(+)/Pi | cotransporter 2C; | | | |
| Entrez Gene | 1426 | 142680 (Human) | | | | |
| SwissProt | Q8N1 | L30 (Human) | | | | |
| Purity | > 97% | 6 | | | | |
| Quality Contro | ol Oligo | nucleotide synthesis | s monitored base by base through trityl analysis to ensure | | | |
| | appro | opriate coupling effici | ency. The oligo is subsequently purified by affinity-solid | | | |
| | phase | e extraction. The anne | ealed RNA duplex is further analyzed by mass | | | |
| | spect | rometry to verify the | exact composition of the duplex. Each lot is compared to | | | |
| | the p | revious lot by mass s | pectrometry to ensure maximum lot-to-lot consistency. | | | |
| Components | We o | ffers pre-designed se | s of 3 different target-specific siRNA oligo duplexes of | | | |
| | huma | an SLC34A3 gene. Eac | h vial contains 5 nmol of lyophilized siRNA. The duplexes | | | |
| | can b | e transfected individu | ally or pooled together to achieve knockdown of the | | | |
| | targe | t gene, which is most | commonly assessed by qPCR or western blot. | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| Component | 15 nmol | 30 nmol |
|---------------------------|--------------|--------------|
| SLC34A3 siRNA (Human) - A | 5 nmol x 1 | 5 nmol x 2 |
| SLC34A3 siRNA (Human) - B | 5 nmol x 1 | 5 nmol x 2 |
| SLC34A3 siRNA (Human) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 μl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 µl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| | | 10 nM | 0.25 μl | 1 μΙ |
| | | 100 nM | 5 µl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| 6-well | 2 ml | 100 nM | 10 µl | 5 µl |
| | | 50 nM | 5 µl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com