

Product Data Sheet

FAM81B siRNA (Human)

| Catalog # | Source | Reactivity | | Applications | | |
|---------------|---|---|--------------------|---------------------|-----------------------|--|
| CRJ5775 | Synthetic | н | | RNAi | | |
| Description | siRNA | to inhibit FAM81B e | xpression using R | NA interference | | |
| Specificity | FAM8 | FAM81B siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed | | | | |
| | to kno | ock down gene expre | ssion. | | | |
| Form | Lyoph | nilized powder | | | | |
| Gene Symbol | FAM8 | FAM81B | | | | |
| Alternative N | ames Prote | in FAM81B | | | | |
| Entrez Gene | 15364 | 43 (Human) | | | | |
| SwissProt | Q96LI | P2 (Human) | | | | |
| Purity | > 97% | 0 | | | | |
| Quality Contr | ol Oligor | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure | | | | |
| | appro | priate coupling effici | ency. The oligo is | subsequently purif | ied by affinity-solid | |
| | phase | e extraction. The anne | ealed RNA duplex | is further analyzed | by mass | |
| | spect | spectrometry to verify the exact composition of the duplex. Each lot is compared to | | | | |
| | the pi | revious lot by mass sp | pectrometry to en | sure maximum lot- | -to-lot consistency. | |
| Components | We of | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of | | | | |
| | huma | human FAM81B gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes | | | | |
| | can b | e transfected individu | ally or pooled to | gether to achieve k | nockdown of the | |
| | target gene, which is most commonly assessed by qPCR or western blot. | | | tern blot. | | |
| | Com | ponent | 1 | 5 nmol | 30 nmol | |
| | FAM | 81B siRNA (Human) - | A 5 | nmol x 1 | 5 nmol x 2 | |
| | FAM | 81B siRNA (Human) - | B 5 | nmol x 1 | 5 nmol x 2 | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| FAM81B siRNA (Human) - C | 5 nmol x 1 | 5 nmol x 2 | |
|--------------------------|--------------|--------------|--|
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 | |
| DEPC Water | 1 ml x 1 | 1 ml x 2 | |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 µl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 µl |
| | | 10 nM | 0.25 μl | 1 µl |
| | | 100 nM | 5 μl | 2 µl |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 µl |
| | | 10 nM | 0.5 μl | 2 µl |
| | | 100 nM | 10 µl | 5 µl |
| 6-well | 2 ml | 50 nM | 5 μl | 5 µl |
| | | 10 nM | 1 µl | 5 µl |

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |