

Product Data Sheet

AWAT1 siRNA (Human)

| Catalog # | Source | Reactivity | Applications | | |
|---------------|-----------|---|-------------------------------------|--------------------------|--|
| CRJ5878 | Synthetic | н | RNAi | | |
| Description | siRNA | to inhibit AWAT1 ex | pression using RNA interference | | |
| Specificity | AWAT | AWAT1 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to | | | |
| | knock | k down gene expressi | on. | | |
| Form | Lyoph | nilized powder | | | |
| Gene Symbol | AWAT | AWAT1 | | | |
| Alternative N | ames DGA2 | DGA2; DGAT2L3; Acyl-CoA wax alcohol acyltransferase 1; Diacylglycerol | | | |
| | O-acy | O-acyltransferase 2-like protein 3; Diacylglycerol acyltransferase 2; | | | |
| | Long- | chain-alcohol O-fatty | -acyltransferase 1 | | |
| Entrez Gene | 15883 | 158833 (Human) | | | |
| SwissProt | Q58H | Q58HT5 (Human) | | | |
| Purity | > 97% | > 97% | | | |
| Quality Contr | ol Oligo | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure | | | |
| | appro | opriate coupling effici | ency. The oligo is subsequently pu | rified by affinity-solid | |
| | phase | e extraction. The anne | ealed RNA duplex is further analyze | ed by mass | |
| | spect | rometry to verify the | exact composition of the duplex. E | Each lot is compared to | |
| | the p | revious lot by mass s | pectrometry to ensure maximum lo | ot-to-lot consistency. | |
| Components | We of | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of | | | |
| | huma | an AWAT1 gene. Each | vial contains 5 nmol of lyophilized | siRNA. The duplexes | |
| | can b | e transfected individu | ally or pooled together to achieve | knockdown of the | |
| | targe | t gene, which is most | commonly assessed by qPCR or we | estern blot. | |
| | Com | ponent | 15 nmol | 30 nmol | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| AWAT1 siRNA (Human) - A | 5 nmol x 1 | 5 nmol x 2 |
|-------------------------|--------------|--------------|
| AWAT1 siRNA (Human) - B | 5 nmol x 1 | 5 nmol x 2 |
| AWAT1 siRNA (Human) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|-----------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 µl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| | | 10 nM | 0.25 μl | 1 μΙ |
| | | 100 nM | 5 μl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| | | 100 nM | 10 µl | 5 µl |
| 6-well | 2 ml | 50 nM | 5 μl | 5 μl |
| . <u></u> | | 10 nM | 1 μl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



For research purposes only, not for human use

Product Data Sheet

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com