

Product Data Sheet

TMEM229B siRNA (Human)

| Reactivity | Applications | | | | |
|---|---|--|---|--|--|
| tic H | RNAi | | | | |
| Description siRNA to inhibit TMEM229B expression using RNA interference | | | | | |
| TMEM229B siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes | | | | | |
| designed to knock down gene expre | ssion. | | | | |
| Lyophilized powder | | | | | |
| TMEM229B | TMEM229B | | | | |
| Iternative Names C14orf83; Transmembrane protein 229B | | | | | |
| Entrez Gene 161145 (Human) | | | | | |
| wissProt Q8NBD8 (Human) | | | | | |
| > 97% | | | | | |
| Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure | | | | | |
| appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid phase extraction. The annealed RNA duplex is further analyzed by mass spectrometry to verify the exact composition of the duplex. Each lot is compared to | | | | | |
| | | | the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency. | | |
| | | | we offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of human TMEM229B gene. Each vial contains 5 nmol of lyophilized siRNA. The | | |
| duplexes can be transfected individually or pooled together to achieve knockdown | | | | | |
| of the target gene, which is most commonly assessed by qPCR or western blot. | | | | | |
| Component | 15 nmol | 30 nmol | | | |
| TMEM229B siRNA (Human) - A | 5 nmol x 1 | 5 nmol x 2 | | | |
| TMEM229B siRNA (Human) - B | 5 nmol x 1 | 5 nmol x 2 | | | |
| | tic H siRNA to inhibit TMEM229B expression TMEM229B siRNA (Human) is a target designed to knock down gene express Lyophilized powder TMEM229B C14orf83; Transmembrane protein 2 161145 (Human) Q8NBD8 (Human) > 97% Oligonucleotide synthesis is monitor appropriate coupling efficiency. The phase extraction. The annealed RNA spectrometry to verify the exact corr the previous lot by mass spectromete We offers pre-designed sets of 3 diff human TMEM229B gene. Each vial co duplexes can be transfected individu of the target gene, which is most corr Component TMEM229B siRNA (Human) - A | tic H RNAi siRNA to inhibit TMEM229B expression using RNA interference TMEM229B siRNA (Human) is a target-specific 19-23 nt siRNA designed to knock down gene expression. Lyophilized powder TMEM229B C14orf83; Transmembrane protein 229B 161145 (Human) Q8NBD8 (Human) > 97% Oligonucleotide synthesis is monitored base by base through appropriate coupling efficiency. The oligo is subsequently pur phase extraction. The annealed RNA duplex is further analyze spectrometry to verify the exact composition of the duplex. E the previous lot by mass spectrometry to ensure maximum lo We offers pre-designed sets of 3 different target-specific siRN human TMEM229B gene. Each vial contains 5 nmol of lyophili duplexes can be transfected individually or pooled together to of the target gene, which is most commonly assessed by qPCF <u>Component</u> 15 nmol TMEM229B siRNA (Human) - A 5 nmol x 1 | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| TMEM229B siRNA (Human) - C | 5 nmol x 1 | 5 nmol x 2 |
|----------------------------|--------------|--------------|
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| 96-well | | 100 nM | 0.5 μl | 0.25 μl |
| | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| 24-well 500 µ | | 100 nM | 2.5 μl | 1 µl |
| | 500 μl | 50 nM | 1.25 μl | 1 µl |
| | | 10 nM | 0.25 μl | 1 µl |
| | | 100 nM | 5 μl | 2 µl |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 µl |
| | | 10 nM | 0.5 μl | 2 µl |
| 6-well | | 100 nM | 10 µl | 5 µl |
| | 2 ml | 50 nM | 5 μΙ | 5 µl |
| | | 10 nM | 1 µl | 5 µl |

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |