

Product Data Sheet

FAM43B siRNA (Human)

Catalog #	Source	Reactivity		Applications		
CRJ6004	Synthetic	н		RNAi		
Description	siRNA	to inhibit FAM43B ex	pression using RI	NA interference		
Specificity	FAM4	3B siRNA (Human) is	a target-specific 1	19-23 nt siRNA oligo	o duplexes designed	
	to kno	ock down gene expre	ssion.			
Form	Lyoph	ilized powder				
Gene Symbol	FAM4	FAM43B				
Alternative N	ames Protei	Protein FAM43B				
Entrez Gene	16393	3 (Human)				
SwissProt	Q6ZT5	52 (Human)				
Purity	> 97%	,				
Quality Contr	ol Oligor	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure				
	appro	priate coupling efficion	ency. The oligo is	subsequently purif	ied by affinity-solid	
	phase	extraction. The anne	ealed RNA duplex	is further analyzed	by mass	
	spectr	rometry to verify the	exact compositio	n of the duplex. Ea	ch lot is compared to	
	the pr	evious lot by mass sp	ectrometry to en	nsure maximum lot-	-to-lot consistency.	
Components We offers pre-designed sets of 3 different target-specific siRNA olig			oligo duplexes of			
	huma	human FAM43B gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes				
	can be	e transfected individu	ally or pooled tog	gether to achieve k	nockdown of the	
	target gene, which is most commonly assessed by qPCR or western blot.			tern blot.		
	Com	ponent	1!	5 nmol	30 nmol	
	FAM	43B siRNA (Human) -	A 5	nmol x 1	5 nmol x 2	
	FAM	43B siRNA (Human) -	B 5	nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

FA	M43B siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Νε	egative Control	2.5 nmol x 1	2.5 nmol x 2
DE	PC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
of medium	of siRNA		2000
	100 nM	0.5 μl	0.25 μl
100 µl	50 nM	0.25 μl	0.25 μl
	10 nM	0.05 μl	0.25 μl
	100 nM	2.5 μl	1 µl
500 μl	50 nM	1.25 μl	1 µl
	10 nM	0.25 μl	1 µl
	100 nM	5 µl	2 µl
1 ml	50 nM	2.5 μl	2 µl
	10 nM	0.5 μl	2 µl
	100 nM	10 µl	5 µl
2 ml	50 nM	5 µl	5 µl
	10 nM	1 µl	5 μl
	of medium 100 μl 500 μl 1 ml	of medium of siRNA 100 nM 100 nM 100 nM 10 nM 50 nM 10 nM 500 μl 50 nM 100 nM 10 nM 500 μl 50 nM 10 nM 10 nM 10 nM 10 nM 10 nM 10 nM 10 nM 10 nM 1 nn 50 nM 1 nn 50 nM 10 nM 50 nM	100 nM0.5 μl100 μl50 nM0.25 μl10 nM0.05 μl100 nM2.5 μl500 μl50 nM1.25 μl10 nM0.25 μl10 nM50 μl100 nM5 μl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com