

Product Data Sheet

LCLAT1 siRNA (Human)

| Catalog # | Source | Reactivity | Applications | | |
|---------------|-----------|---|---|--------------------|--|
| CRJ6644 | Synthetic | Н | RNAi | | |
| Description | siRNA | A to inhibit LCLAT1 exp | pression using RNA interference | | |
| Specificity | LCLAT | LCLAT1 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to | | | |
| | knocl | k down gene expressi | on. | | |
| Form | Lyoph | nilized powder | | | |
| Gene Symbol | LCLAT | LCLAT1 | | | |
| Alternative N | ames AGPA | AGPAT8; ALCAT1; LYCAT; Lysocardiolipin acyltransferase 1; | | | |
| | 1-acy | lglycerol-3-phosphate | e O-acyltransferase 8; 1-AGP acyltransfer | rase 8; 1-AGPAT 8; | |
| | Acyl-0 | CoA:lysocardiolipin ac | yltransferase 1 | | |
| Entrez Gene | 2535 | 58 (Human) | | | |
| SwissProt | Q6UV | Q6UWP7 (Human) | | | |
| Purity | > 97% | > 97% | | | |
| Quality Contr | ol Oligo | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure | | | |
| | appro | opriate coupling effici | ency. The oligo is subsequently purified | by affinity-solid | |
| | phase | e extraction. The anne | ealed RNA duplex is further analyzed by | mass | |
| | spect | rometry to verify the | exact composition of the duplex. Each lo | ot is compared to | |
| | the p | revious lot by mass sp | pectrometry to ensure maximum lot-to-l | ot consistency. | |
| Components | We o | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of | | | |
| | huma | an LCLAT1 gene. Each | vial contains 5 nmol of lyophilized siRNA | . The duplexes | |
| | can b | e transfected individu | ally or pooled together to achieve knoc | kdown of the | |
| | targe | target gene, which is most commonly assessed by qPCR or western blot. | | | |
| | Com | ponent | 15 nmol 30 | nmol | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| LCLAT1 siRNA (Human) - A | 5 nmol x 1 | 5 nmol x 2 |
|--------------------------|--------------|--------------|
| LCLAT1 siRNA (Human) - B | 5 nmol x 1 | 5 nmol x 2 |
| LCLAT1 siRNA (Human) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μΙ |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μl |
| | | 10 nM | 0.25 μl | 1 μΙ |
| | | 100 nM | 5 µl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| | | 100 nM | 10 µl | 5 µl |
| 6-well | 2 ml | 50 nM | 5 μl | 5 μl |
| | | 10 nM | 1 μl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



For research purposes only, not for human use

Product Data Sheet

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com