

# **Product Data Sheet**

### **OR13H1 siRNA (Human)**

| Catalog #     | Source      | Reactivity  |               | Applications           |                        |  |
|---------------|-------------|---|---------------|------------------------|------------------------|--|
| CRJ7504       | Synthetic   | ,<br>H  |               | RNAi                   |                        |  |
| Description   | siRNA       | siRNA to inhibit OR13H1 expression using RNA interference                             |               |                        |                        |  |
| Specificity   | OR13I       | OR13H1 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed      |               |                        |                        |  |
|               | to kno      | ock down gene expres  | ssion.        |                        |                        |  |
| Form          | Lyoph       | ilized powder   |               |                        |                        |  |
| Gene Symbol   | OR13I       | OR13H1  |               |                        |                        |  |
| Alternative N | ames Olfact | Olfactory receptor 13H1; Olfactory receptor ORX-1                                     |               |                        |                        |  |
| Entrez Gene   | 34746       | 68 (Human)  |               |                        |                        |  |
| SwissProt     | Q8NG        | 92 (Human)  |               |                        |                        |  |
| Purity        | > 97%       | > 97%   |               |                        |                        |  |
| Quality Contr | ol Oligor   | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure |               |                        |                        |  |
|               | appro       | appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid |               |                        |                        |  |
|               | phase       | extraction. The anne  | aled RNA dup  | lex is further analyze | ed by mass             |  |
|               | spectr      | spectrometry to verify the exact composition of the duplex. Each lot is compared to   |               |                        |                        |  |
|               | the pr      | evious lot by mass sp   | ectrometry to | ensure maximum lo      | ot-to-lot consistency. |  |
| Components    | We of       | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of    |               |                        |                        |  |
|               | huma        | human OR13H1 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes       |               |                        |                        |  |
|               | can be      | can be transfected individually or pooled together to achieve knockdown of the        |               |                        |                        |  |
|               | target      | target gene, which is most commonly assessed by qPCR or western blot.                 |               |                        |                        |  |
|               | Com         | ponent  |               | 15 nmol                | 30 nmol                |  |
|               | OR13        | 3H1 siRNA (Human) -   | A             | 5 nmol x 1             | 5 nmol x 2             |  |
|               | OR13        | 3H1 siRNA (Human) -   | В             | 5 nmol x 1             | 5 nmol x 2             |  |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



## **Product Data Sheet**

| DEPC Water               | 1 ml x 1     | 1 ml x 2     |
|--------------------------|--------------|--------------|
| Negative Control         | 2.5 nmol x 1 | 2.5 nmol x 2 |
| OR13H1 siRNA (Human) - C | 5 nmol x 1   | 5 nmol x 2   |

**Directions for Use** We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to

cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

| Plate   | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
|         | of medium    | of siRNA            |               | 2000         |
|         |              | 100 nM              | 0.5 μl        | 0.25 μl      |
| 96-well | 100 µl       | 50 nM               | 0.25 μl       | 0.25 μl      |
|         |              | 10 nM               | 0.05 μl       | 0.25 μl      |
|         |              | 100 nM              | 2.5 μl        | 1 µl         |
| 24-well | 500 μl       | 50 nM               | 1.25 μl       | 1 µl         |
|         |              | 10 nM               | 0.25 μl       | 1 µl         |
|         |              | 100 nM              | 5 μl          | 2 µl         |
| 12-well | 1 ml         | 50 nM               | 2.5 μl        | 2 µl         |
|         |              | 10 nM               | 0.5 μl        | 2 µl         |
|         |              | 100 nM              | 10 µl         | 5 µl         |
| 6-well  | 2 ml         | 50 nM               | 5 μl          | 5 μl         |
|         |              | 10 nM               | 1 µl          | 5 μl         |

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |