

# **Product Data Sheet**

## NMNAT3 siRNA (Human)

ource	Reactivity	Applications		
-		RNAi		
siRNA	siRNA to inhibit NMNAT3 expression using RNA interference			
NMNA	NMNAT3 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed			
to kno	ck down gene expres	sion.		
Lyophi	ilized powder	zed powder		
NMNA	AT3			
es Nicotir	Nicotinamide mononucleotide adenylyltransferase 3; NMN adenylyltransferase 3;			
Nicotir	nate-nucleotide aden	ylyltransferase 3; NaMN adenylyltransferase 3; Pyridine		
nuclec	otide adenylyltransfer	ase 3; PNAT-3		
34956	5 (Human)			
Q96T6				
> 97%	> 97%			
Oligon	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure			
approp	priate coupling efficie	ency. The oligo is subsequently purified by affinity-solid		
phase	extraction. The anne	aled RNA duplex is further analyzed by mass		
spectre	ometry to verify the	exact composition of the duplex. Each lot is compared to		
the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.				
We off	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
human NMNAT3 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes				
can be	e transfected individu	ally or pooled together to achieve knockdown of the		
target gene, which is most commonly assessed by qPCR or western blot.				
Comp	ponent	15 nmol 30 nmol		
	NMNA to kno Lyophi NMNA es Nicotii Nicotii nucleo 34956 Q96T6 > 97% Oligon appro phase spectr the pr We off human can be target	ynthetic H siRNA to inhibit NMNAT3 ex NMNAT3 siRNA (Human) is to knock down gene express Lyophilized powder NMNAT3 es Nicotinamide mononucleot Nicotinate-nucleotide aden nucleotide adenylyltransfer 349565 (Human) Q96T66 (Human) > 97% Oligonucleotide synthesis is appropriate coupling efficie phase extraction. The anne spectrometry to verify the o the previous lot by mass sp We offers pre-designed sets human NMNAT3 gene. Each can be transfected individu		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

## **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com

# **Product Data Sheet**

NMNAT3 siRNA (Human) - A	5 nmol x 1	5 nmol x 2
NMNAT3 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
NMNAT3 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μΙ
24-well	500 μl	50 nM	1.25 μl	1 μl
		10 nM	0.25 μl	1 μΙ
		100 nM	5 µl	2 μΙ
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 μl
		10 nM	1 μl	5 μΙ

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



For research purposes only, not for human use

## **Product Data Sheet**

**Storage/Stability** Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com