

Product Data Sheet

GTF2IRD2B siRNA (Human)

Catalog #	Source	Reactivity	Applications		
CRJ7887	Synthetic	н	RNAi		
Description	siRNA	siRNA to inhibit GTF2IRD2B expression using RNA interference			
Specificity	GTF2I	RD2B siRNA (Human)	is a target-specific 19-23 nt siRl	NA oligo duplexes	
	desigr	designed to knock down gene expression.			
Form	Lyoph	Lyophilized powder			
Gene Symbol	GTF2I	GTF2IRD2B			
Alternative N	ames Gener	General transcription factor II-I repeat domain-containing protein 2B; GTF2I repeat			
	domai	in-containing protein	2B; Transcription factor GTF2IR	D2-beta	
Entrez Gene	38952	389524 (Human)			
SwissProt	Q6EKJ	Q6EKJ0 (Human)			
Purity	> 97%	> 97%			
Quality Contr	ol Oligor	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure			
	appro	priate coupling efficie	ncy. The oligo is subsequently p	ourified by affinity-solid	
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass			
	spectr	spectrometry to verify the exact composition of the duplex. Each lot is compared to			
	the pr	the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.			
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	huma	human GTF2IRD2B gene. Each vial contains 5 nmol of lyophilized siRNA. The			
	duplex	duplexes can be transfected individually or pooled together to achieve knockdown			
	of the	of the target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	GTF2	IRD2B siRNA (Human) - A 5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

DEPC Water	1 ml x 1	1 ml x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
GTF2IRD2B siRNA (Human) - C	5 nmol x 1	5 nmol x 2
GTF2IRD2B siRNA (Human) - B	5 nmol x 1	5 nmol x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 μl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
		100 nM	10 µl	5 μΙ
6-well	2 ml	50 nM	5 µl	5 μΙ
		10 nM	1 μΙ	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com