

# **Product Data Sheet**

## USP27X siRNA (Human)

ource	Reactivity	Applications		
nthetic/	н	RNAi		
siRNA	to inhibit USP27X ex	pression using RNA interference		
USP27	USP27X siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to			
knock	down gene expressio	on.		
Lyoph	ilized powder	ized powder		
Gene Symbol USP27				
es USP22	USP22L; USP27; Ubiquitin carboxyl-terminal hydrolase 27; Deubiquitinating enzyme			
27; Ut	piquitin carboxyl-tern	ninal hydrolase 22-like; Ubiquitin thioesterase 27;		
Ubiqu	itin-specific-processi	ng protease 27; X-linked ubiquitin carboxyl-terminal		
hydro	lase 27			
38985	389856 (Human)			
A6NN	A6NNY8 (Human)			
> 97%	> 97%			
Oligor	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure			
appro	appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid			
phase	phase extraction. The annealed RNA duplex is further analyzed by mass			
spectr	rometry to verify the	exact composition of the duplex. Each lot is compared to		
the pr	evious lot by mass sp	ectrometry to ensure maximum lot-to-lot consistency.		
We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
huma	n USP27X gene. Each	vial contains 5 nmol of lyophilized siRNA. The duplexes		
can be	e transfected individu	ally or pooled together to achieve knockdown of the		
target	gene, which is most	commonly assessed by qPCR or western blot.		
	vnthetic siRNA USP27 knock Lyoph USP27 27; Ut Ubiqu hydro 38985 A6NN > 97% Oligor appro phase spectr the pr We of huma can be	Anthetic     H       siRNA to inhibit USP27X ex       USP27X siRNA (Human) is a       knock down gene expression       Lyophilized powder       USP27X       Sologen (Lyophilized powder)       USP27X       USP27X       Sologen (Lyophilized powder)       USP27X       Sologen (Lyophilized powder)       USP27X       Sologen (Lyophilized powder)       Ubiquitin-specific-processing       hydrolase 27       389856 (Human)       > 97%       Oligonucleotide synthesis i       appropriate coupling efficiend       phase extraction. The anneed       spectrometry to verify the       the previous lot by mass spectrometry to verify the       human USP27X gene. Each       can be transfected individue		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

Component	15 nmol	30 nmol
USP27X siRNA (Human) - A	5 nmol x 1	5 nmol x 2
USP27X siRNA (Human) - B	5 nmol x 1	5 nmol x 2
USP27X siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 μl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μl
		10 nM	0.25 μl	1 μl
		100 nM	5 µl	2 μΙ
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
6-well	2 ml	100 nM	10 µl	5 µl
		50 nM	5 µl	5 μΙ

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com