

# **Product Data Sheet**

### KRTAP5-11 siRNA (Human)

Catalog #	Source	Reactivity	Applications		
CRJ8230	Synthetic	н	RNAi		
Description	siRNA	to inhibit KRTAP5-1	expression using RNA interferen	ice	
Specificity	KRTAI	P5-11 siRNA (Human)	is a target-specific 19-23 nt siRN	A oligo duplexes	
	desig	ned to knock down g	ene expression.		
Form	Lyoph	nilized powder			
Gene Symbol	KRTAI	KRTAP5-11			
Alternative N	ames KAP5	KAP5.11; KRTAP5.11; Keratin-associated protein 5-11; Keratin-associated protein			
	5.11;	Ultrahigh sulfur kera	tin-associated protein 5.11		
Entrez Gene	4400	51 (Human)			
SwissProt	Q6L8	Q6L8G4 (Human)			
Purity	> 97%	> 97%			
Quality ControlOligonucleotide synthesis is monitored base by			is monitored base by base throug	trityl analysis to ensure	
	appro	opriate coupling effici	ency. The oligo is subsequently p	urified by affinity-solid	
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass			
	spect	spectrometry to verify the exact composition of the duplex. Each lot is compared to			
	the p	revious lot by mass s	pectrometry to ensure maximum	lot-to-lot consistency.	
Components	We o	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	huma	human KRTAP5-11 gene. Each vial contains 5 nmol of lyophilized siRNA. The			
	duple	duplexes can be transfected individually or pooled together to achieve knockdown			
	of the	of the target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	KRTA	AP5-11 siRNA (Humar	n) - A 5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

KRTAP5-11 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
KRTAP5-11 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 μl
		10 nM	1 μl	5 μl

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com