

# **Product Data Sheet**

## DMRTC1B siRNA (Human)

Catalog #	Source	Reactivity		Applications		
CRJ8686	Synthetic	н		RNAi		
Description	siRNA	siRNA to inhibit DMRTC1B expression using RNA interference				
Specificity	DMR	DMRTC1B siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed				
	to kno	ock down gene expres	ssion.			
Form	Lyoph	Lyophilized powder				
Gene Symbol	DMR	DMRTC1B				
Alternative N	ames Doub	Doublesex- and mab-3-related transcription factor C1				
Entrez Gene	7286	728656 (Human)				
SwissProt	Q5HY	Q5HYR2 (Human)				
Purity	> 97%	> 97%				
Quality Control Oligonucleotide synthesis is monitored base by base through trityl and			trityl analysis to ensure			
	appro	priate coupling efficie	ency. The oligo	is subsequently pur	ified by affinity-solid	
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass				
	spect	spectrometry to verify the exact composition of the duplex. Each lot is compared to				
	the p	revious lot by mass sp	ectrometry to	ensure maximum lo	t-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of				
	huma	in DMRTC1B gene. Ea	ch vial contains	5 nmol of lyophilize	ed siRNA. The duplexes	
	can b	e transfected individu	ally or pooled t	together to achieve	knockdown of the	
	target	target gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent		15 nmol	30 nmol	
	DMF	RTC1B siRNA (Human)	- A	5 nmol x 1	5 nmol x 2	
	D1 45		-			

DMRTC1B siRNA (Human) - B5 nmol x 15 nmol x 2Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-<br/>Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility<br/>Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

DMRTC1B siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 μl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 µl

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com