

Product Data Sheet

ADRB2 siRNA (Mouse)

Catalog #	Source	Reactivity	Δηρί	ications	
-					
CRM0090	Synthetic	Μ	RNAi		
Description	siRNA	to inhibit ADRB2 exp	pression using RNA inte	erference	
Specificity	ADRB	2 siRNA (Mouse) is a	target-specific 19-23 n	t siRNA oligo duplexes designed to	
	knock	down gene expressi	on.		
Form	Lyoph	ilized powder			
Gene Symbol	ADRB	2			
Alternative N	ames ADRB	2R; Beta-2 adrenergi	c receptor; Beta-2 adre	enoreceptor; Beta-2 adrenoceptor	
Entrez Gene	11555	5 (Mouse)			
SwissProt	P1876	62 (Mouse)			
Purity	> 97%	, D			
Quality Control Oligonucleotide synthesis is monitored base by base through trityl analys			ase through trityl analysis to ensure		
	appro	priate coupling effici	ency. The oligo is subse	equently purified by affinity-solid	
	phase	extraction. The anne	ealed RNA duplex is fur	ther analyzed by mass	
	spect	rometry to verify the	exact composition of t	he duplex. Each lot is compared to	
	the pi	revious lot by mass sp	bectrometry to ensure	maximum lot-to-lot consistency.	
Components	We of	ffers pre-designed set	s of 3 different target-	specific siRNA oligo duplexes of	
	mous	mouse ADRB2 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes			
	can b	can be transfected individually or pooled together to achieve knockdown of the			
	target	target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nm	ol 30 nmol	
	ADR	B2 siRNA (Mouse) - A	5 nmo	l x 1 5 nmol x 2	
	ADR	B2 siRNA (Mouse) - B	5 nmo	l x 1 5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

Negative Control2.5 nmol x 12.5 nmol x 2DEPC Water1 ml x 11 ml x 2	25 nmol x 1 25 nmol x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com