

Product Data Sheet

ATP6V0E1 siRNA (Mouse)

Catalog #	Source	Reactivity	Applic	ations	
CRM0312	Synthetic	М	RNAi		
Description	siRNA	to inhibit ATP6V0E1	expression using RNA in	terference	
Specificity	ATP6	V0E1 siRNA (Mouse) i	s a target-specific 19-23	nt siRNA oligo duplexes designed	
	to kno	ock down gene expre	ssion.		
Form	Lyoph	nilized powder			
Gene Symbol	ATP6	ATP6V0E1			
Alternative N	ames ATP6	ATP6V0E; V-type proton ATPase subunit e 1; V-ATPase subunit e 1; Vacuolar proton			
	pump	o subunit e 1			
Entrez Gene	11974	4 (Mouse)			
SwissProt	Q9CC	Q9CQD8 (Mouse)			
Purity	> 97%	> 97%			
Quality Contr	ol Oligo	Digonucleotide synthesis is monitored base by base through trityl analysis to ensure			
	appro	opriate coupling effici	ency. The oligo is subsec	uently purified by affinity-solid	
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass			
	spect	spectrometry to verify the exact composition of the duplex. Each lot is compared to			
	the p	the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.			
Components	We o	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	mous	mouse ATP6V0E1 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes			
	can b	can be transfected individually or pooled together to achieve knockdown of the			
	targe	target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	ATP	6V0E1 siRNA (Mouse)	- A 5 nmol >	1 5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

ATP6V0E1 siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
ATP6V0E1 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
_		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
_		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 µl	5 μΙ
		10 nM	1 μΙ	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com