

Product Data Sheet

SLC7A3 siRNA (Mouse)

Catalog #	Source	Reactivity	Applications			
CRM0323	Synthetic	М	RNAi			
Description	siRNA	to inhibit SLC7A3 ex	pression using RNA interference			
Specificity	SLC7A	SLC7A3 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed to				
	knock	down gene express	on.			
Form	Lyoph	nilized powder				
Gene Symbol	SLC7A	SLC7A3				
Alternative N	ames ATRC	ATRC3; CAT3; Cationic amino acid transporter 3; CAT-3; CAT3; Cationic amino acid				
	transp	porter y+; Solute car	ier family 7 member 3			
Entrez Gene	11989	9 (Mouse)				
SwissProt	P7042	P70423 (Mouse)				
Purity	> 97%	> 97%				
Quality Control Oligonucleotide synthesis is monitored base by base through trityl analy			h trityl analysis to ensure			
	appro	appropriate coupling efficiency. The oligo is subsequently purified by affinity-soli				
	phase	e extraction. The ann	ealed RNA duplex is further analyz	zed by mass		
	spect	rometry to verify the	exact composition of the duplex.	Each lot is compared to		
	the p	revious lot by mass s	pectrometry to ensure maximum	lot-to-lot consistency.		
Components We offers pre-designed sets of 3 different target-specific siRNA oligo duplexe				NA oligo duplexes of		
	mous	e SLC7A3 gene. Each	vial contains 5 nmol of lyophilized	d siRNA. The duplexes		
	can b	e transfected individ	ually or pooled together to achiev	e knockdown of the		
	target	target gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent	15 nmol	30 nmol		
	SLC7	A3 siRNA (Mouse) -	A 5 nmol x 1	5 nmol x 2		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

SLC7A3 siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
SLC7A3 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 μl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
		100 nM	10 µl	5 μΙ
6-well	2 ml	50 nM	5 μl	5 μΙ
		10 nM	1 µl	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com