

# **Product Data Sheet**

## BST1 siRNA (Mouse)

Catalog #	Source	Reactivity	Applications			
CRM0410	Synthetic	М	RNAi			
Description	siRNA	to inhibit BST1 expre	ession using RNA interference			
Specificity	BST1	siRNA (Mouse) is a ta	rget-specific 19-23 nt siRNA oligo duplexes designed to			
	knock	<pre>&lt; down gene expression</pre>	on.			
Form	Lyoph	nilized powder				
Gene Symbol	BST1					
Alternative Na	imes BP-3;	BP-3; BP3; LY65; ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 2; ADP-ribosyl				
	cyclas	se 2; Antigen BP3; BP-	3 alloantigen; Bone marrow stromal antigen 1; BST-1;			
	Cyclic	: ADP-ribose hydrolas	e 2; cADPr hydrolase 2; Leukocyte antigen 65; Ly-65; CD			
	antige	en CD157				
Entrez Gene	12182	2 (Mouse)				
SwissProt	Q642	77 (Mouse)				
Purity	> 97%	6				
Quality Contro	ol Oligo	nucleotide synthesis i	s monitored base by base through trityl analysis to ensure			
	appro	opriate coupling efficion	ency. The oligo is subsequently purified by affinity-solid			
	phase	e extraction. The anne	ealed RNA duplex is further analyzed by mass			
	spect	rometry to verify the	exact composition of the duplex. Each lot is compared to			
	the p	revious lot by mass sp	pectrometry to ensure maximum lot-to-lot consistency.			
Components	We o	ffers pre-designed set	s of 3 different target-specific siRNA oligo duplexes of			
	mous	e BST1 gene. Each via	l contains 5 nmol of lyophilized siRNA. The duplexes can			
	be tra	ansfected individually	or pooled together to achieve knockdown of the target			
	gene,	which is most comm	only assessed by qPCR or western blot.			
		An at a way belief at the law	kistaskanistas II. kunsta fluorena II. E. I.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

Component	15 nmol	30 nmol
BST1 siRNA (Mouse) - A	5 nmol x 1	5 nmol x 2
BST1 siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
BST1 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 μl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μl
		10 nM	0.25 μl	1 μl
		100 nM	5 μl	2 μΙ
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
6-well	2 ml	100 nM	10 µl	5 µl
o-well	2 ml	50 nM	5 μl	5 μΙ

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com