

Product Data Sheet

BTG2 siRNA (Mouse)

| Catalog # | Source | Reactivity | Applications | | |
|--|-------------|---|-----------------------------------|--------------------------|--|
| CRM0424 | Synthetic | Μ | RNAi | | |
| Description | siRNA | to inhibit BTG2 expres | sion using RNA interference | | |
| Specificity | BTG2 s | BTG2 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed to | | | |
| | knock | knock down gene expression. | | | |
| Form | Lyophi | Lyophilized powder | | | |
| Gene Symbol | BTG2 | BTG2 | | | |
| Alternative Na | ames TIS21; | TIS21; Protein BTG2; BTG family member 2; NGF-inducible protein TIS21 | | | |
| Entrez Gene | 12227 | 12227 (Mouse) | | | |
| SwissProt | Q0421 | Q04211 (Mouse) | | | |
| Purity > 97% | | | | | |
| Quality Control Oligonucleotide synthesis is monitored base by base through trityl | | n trityl analysis to ensure | | | |
| | appro | priate coupling efficier | ncy. The oligo is subsequently pu | rified by affinity-solid | |
| | phase | phase extraction. The annealed RNA duplex is further analyzed by mass | | | |
| | spectr | spectrometry to verify the exact composition of the duplex. Each lot is compared to | | | |
| | the pr | evious lot by mass spe | ctrometry to ensure maximum l | ot-to-lot consistency. | |
| Components | We of | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of | | | |
| | mouse | e BTG2 gene. Each vial | contains 5 nmol of lyophilized si | RNA. The duplexes can | |
| | be tra | nsfected individually o | r pooled together to achieve kno | ockdown of the target | |
| gene, which is most commonly assessed by qPCR or western blot. | | | blot. | | |
| | Com | ponent | 15 nmol | 30 nmol | |
| | BTG2 | siRNA (Mouse) - A | 5 nmol x 1 | 5 nmol x 2 | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

5 nmol x 1

5 nmol x 2

BTG2 siRNA (Mouse) - B

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| BTG2 siRNA (Mouse) - C | 5 nmol x 1 | 5 nmol x 2 |
|------------------------|--------------|--------------|
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 µl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 µl |
| | | 10 nM | 0.25 μl | 1 µl |
| | | 100 nM | 5 μl | 2 µl |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 µl |
| | | 10 nM | 0.5 μl | 2 µl |
| | | 100 nM | 10 µl | 5 µl |
| 6-well | 2 ml | 50 nM | 5 μl | 5 µl |
| | | 10 nM | 1 µl | 5 µl |

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |