

Product Data Sheet

DDR1 siRNA (Mouse)

| Catalog # | Source | Reactivity | Applications | | | |
|------------------------|-----------|---|--|--|--|--|
| CRM0461 | Synthetic | Μ | RNAi | | | |
| Description | siRN | A to inhibit DDR1 exp | ession using RNA interference | | | |
| Specificity | DDR | 1 siRNA (Mouse) is a t | arget-specific 19-23 nt siRNA oligo duplexes designed to | | | |
| | knoc | k down gene expressi | on. | | | |
| Form | Lyop | hilized powder | | | | |
| Gene Symbol | DDR: | DDR1 | | | | |
| Alternative Na | ames CAK; | CAK; EDDR1; MPK6; Epithelial discoidin domain-containing receptor 1; Epithelial | | | | |
| | disco | idin domain receptor | 1; CD167 antigen-like family member A; Cell adhesion | | | |
| | kinas | e; Discoidin receptor | tyrosine kinase; Protein-tyrosine kinase MPK-6; Tyrosine | | | |
| | kinas | se DDR; | | | | |
| Entrez Gene | 1230 | 5 (Mouse) | | | | |
| SwissProt | Q031 | L46 (Mouse) | | | | |
| Purity | > 979 | % | | | | |
| Quality Contro | ol Oligo | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure | | | | |
| | appro | opriate coupling effici | ency. The oligo is subsequently purified by affinity-solid | | | |
| | phas | e extraction. The ann | ealed RNA duplex is further analyzed by mass | | | |
| | spect | trometry to verify the | exact composition of the duplex. Each lot is compared to | | | |
| | the p | previous lot by mass s | pectrometry to ensure maximum lot-to-lot consistency. | | | |
| Components | We c | offers pre-designed se | ts of 3 different target-specific siRNA oligo duplexes of | | | |
| | mou | se DDR1 gene. Each v | al contains 5 nmol of lyophilized siRNA. The duplexes can | | | |
| | be tr | ansfected individually | or pooled together to achieve knockdown of the target | | | |
| | gene | , which is most comm | only assessed by qPCR or western blot. | | | |
| A secold secold second | | Alexterne blet 111 because | | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| Component | 15 nmol | 30 nmol |
|------------------------|--------------|--------------|
| DDR1 siRNA (Mouse) - A | 5 nmol x 1 | 5 nmol x 2 |
| DDR1 siRNA (Mouse) - B | 5 nmol x 1 | 5 nmol x 2 |
| DDR1 siRNA (Mouse) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μΙ |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| | | 10 nM | 0.25 μl | 1 μΙ |
| | | 100 nM | 5 µl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| 6-well | 2 ml | 100 nM | 10 µl | 5 µl |
| o-well | z mi | 50 nM | 5 µl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com