

Product Data Sheet

CSNK2A2 siRNA (Mouse)

Catalog #	Source	Reactivity		Applications		
CRM0883	Synthetic	Μ		RNAi		
Description	siRNA	to inhibit CSNK2A2	expression using	g RNA interference		
Specificity	CSNK	CSNK2A2 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed				
	to kno	ock down gene expre	ssion.			
Form	Lyoph	ilized powder				
Gene Symbol	CSNK:	CSNK2A2				
Alternative N	ames Caseii	n kinase II subunit alp	oha'; CK II alpha	1		
Entrez Gene	13000) (Mouse)				
SwissProt	0548	O54833 (Mouse)				
Purity	> 97%	> 97%				
Quality Contr	ol Oligoi	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure				
	appro	appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid				
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass				
	spect	spectrometry to verify the exact composition of the duplex. Each lot is compared to				
	the pi	the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.				
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of				
	mous	mouse CSNK2A2 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes				
	can b	can be transfected individually or pooled together to achieve knockdown of the				
	target	target gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent		15 nmol	30 nmol	
	CSNI	K2A2 siRNA (Mouse)	- A	5 nmol x 1	5 nmol x 2	
	CSNI	K2A2 siRNA (Mouse)	- B	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

DEPC Water	1 ml x 1	1 ml x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
CSNK2A2 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 μl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com