

## **Product Data Sheet**

### CYP1B1 siRNA (Mouse)

Catalog #	Source	Reactivity	Applications		
CRM0931	Synthetic	Μ	RNAi		
Description	siRNA	to inhibit CYP1B1 ex	pression using RNA interference		
Specificity	CYP1E	CYP1B1 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed to			
	knock	down gene expressio	on.		
Form	Lyoph	Lyophilized powder			
Gene Symbol	CYP1E	CYP1B1			
Alternative N	ames CYP1-	CYP1-B1; Cytochrome P450 1B1; CYPIB1; Cytochrome P450CMEF; Cytochrome			
	P450E	F			
Entrez Gene	13078	3 (Mouse)			
SwissProt	Q6442	29 (Mouse)			
Purity > 97%					
Quality Control Oligonucleotide synthesis is monitored base by base through trityl analysis			gh trityl analysis to ensure		
	appro	priate coupling efficie	ency. The oligo is subsequently p	urified by affinity-solid	
	phase	extraction. The anne	aled RNA duplex is further analy	zed by mass	
	spectr	rometry to verify the	exact composition of the duplex.	. Each lot is compared to	
	the pr	revious lot by mass sp	ectrometry to ensure maximum	lot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	mouse	e CYP1B1 gene. Each	vial contains 5 nmol of lyophilize	d siRNA. The duplexes	
	can be	e transfected individu	ally or pooled together to achiev	ve knockdown of the	
	target	target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	CYP1	.B1 siRNA (Mouse) - A	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



# **Product Data Sheet**

CYP1B1 siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
CYP1B1 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 μl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
		100 nM	10 µl	5 μΙ
6-well	2 ml	50 nM	5 μl	5 μΙ
		10 nM	1 µl	5 μΙ

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com