

Product Data Sheet

ECEL1 siRNA (Mouse)

Catalog #	Source	Reactivity	Applications		
CRM1121	Synthetic	Μ	RNAi		
Description	siRNA	to inhibit ECEL1 expr	ession using RNA interference		
Specificity	ECEL1	. siRNA (Mouse) is a t	arget-specific 19-23 nt siRNA oligo	o duplexes designed to	
	knock	down gene expressio	on.		
Form	Lyoph	ilized powder			
Gene Symbol	ECEL1	ECEL1			
Alternative N	ames DINE;	DINE; XCE; Endothelin-converting enzyme-like 1; Damage-induced neuronal			
	endor	peptidase; Xce proteii	ı		
Entrez Gene	13599) (Mouse)			
SwissProt	Q9JM	I0 (Mouse)			
Purity	> 97%	> 97%			
Quality Control Oligonucleotide synthesis is m			s monitored base by base through	n trityl analysis to ensure	
	appro	priate coupling efficie	ency. The oligo is subsequently pu	rified by affinity-solid	
	phase	extraction. The anne	aled RNA duplex is further analyz	ed by mass	
	spect	rometry to verify the	exact composition of the duplex.	Each lot is compared to	
	the pr	revious lot by mass sp	ectrometry to ensure maximum l	ot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	mous	e ECEL1 gene. Each vi	al contains 5 nmol of lyophilized s	siRNA. The duplexes can	
	be tra	be transfected individually or pooled together to achieve knockdown of the target			
	gene,	gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	ECEL	1 siRNA (Mouse) - A	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

ECEL1 siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
ECEL1 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μl
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 μl
12-well	1 ml	50 nM	2.5 μl	2 μl
_		10 nM	0.5 μl	2 μl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com