

Product Data Sheet

GOT2 siRNA (Mouse)

| Source | Reactivity | Applications | | |
|-----------|---|---|--|--|
| Synthetic | Μ | RNAi | | |
| siRN | A to inhibit GOT2 exp | ression using RNA interference | | |
| GOT2 | GOT2 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed to | | | |
| knoc | k down gene expressi | on. | | |
| Lyop | hilized powder | | | |
| GOT2 | GOT2 | | | |
| mes GOT- | GOT-2; Aspartate aminotransferase mitochondrial; mAspAT; Fatty acid-binding | | | |
| prote | ein; FABP-1; Glutamat | e oxaloacetate transaminase 2; Kynurenine | | |
| amin | otransferase 4; Kynur | enine aminotransferase IV; Kynurenineoxoglutarate | | |
| trans | aminase 4; Kynurenir | neoxoglut | | |
| 1471 | 14719 (Mouse) | | | |
| P052 | P05202 (Mouse) | | | |
| > 97% | 6 | | | |
| l Oligo | nucleotide synthesis | is monitored base by base through trityl analysis to ensure | | |
| appro | opriate coupling effici | ency. The oligo is subsequently purified by affinity-solid | | |
| phas | e extraction. The ann | ealed RNA duplex is further analyzed by mass | | |
| spect | trometry to verify the | exact composition of the duplex. Each lot is compared to | | |
| the p | revious lot by mass s | pectrometry to ensure maximum lot-to-lot consistency. | | |
| We o | ffers pre-designed se | ts of 3 different target-specific siRNA oligo duplexes of | | |
| mous | se GOT2 gene. Each v | ial contains 5 nmol of lyophilized siRNA. The duplexes can | | |
| be tr | ansfected individually | or pooled together to achieve knockdown of the target | | |
| gene | , which is most comm | only assessed by qPCR or western blot. | | |
| | Synthetic siRN/ GOT2 knoc Lyop GOT2 mes GOT- prote amin trans 1471 P052 > 979 I Oligo appro phas spect the p We o mous | Synthetic M siRNA to inhibit GOT2 expl GOT2 siRNA (Mouse) is a to knock down gene expressi Lyophilized powder GOT2 mes GOT-2; Aspartate aminotrate protein; FABP-1; Glutamat aminotransferase 4; Kynure transaminase 4; Kynurenir 14719 (Mouse) P05202 (Mouse) > 97% Oligonucleotide synthesis appropriate coupling effici phase extraction. The anne spectrometry to verify the the previous lot by mass s We offers pre-designed see mouse GOT2 gene. Each vi be transfected individually | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| Component | 15 nmol | 30 nmol |
|------------------------|--------------|--------------|
| GOT2 siRNA (Mouse) - A | 5 nmol x 1 | 5 nmol x 2 |
| GOT2 siRNA (Mouse) - B | 5 nmol x 1 | 5 nmol x 2 |
| GOT2 siRNA (Mouse) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| 96-well | 100 µl | 100 nM | 0.5 μl | 0.25 μl |
| | | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μΙ |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| | | 10 nM | 0.25 μl | 1 μl |
| | | 100 nM | 5 μl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| 6-well | 2 ml | 100 nM | 10 µl | 5 μΙ |
| | | 50 nM | 5 µl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com