

Product Data Sheet

FOXJ1 siRNA (Mouse)

Catalog #	Source	Reactivity	Applications		
CRM1817	Synthetic	М	RNAi		
Description	siRNA	to inhibit FOXJ1 exp	ression using RNA interference		
Specificity	FOXJ1	L siRNA (Mouse) is a t	arget-specific 19-23 nt siRNA oligo	oduplexes designed to	
	knock	down gene expressi	on.		
Form	Lyoph	nilized powder			
Gene Symbol	FOXJ1	FOXJ1			
Alternative N	ames FKHL1	FKHL13; HFH4; Forkhead box protein J1; Forkhead-related protein FKHL13;			
	Hepat	tocyte nuclear factor	3 forkhead homolog 4; HFH-4		
Entrez Gene	15223	3 (Mouse)			
SwissProt	Q616	Q61660 (Mouse)			
Purity > 97%		97%			
Quality Contr	y Control Oligonucleotide synthesis is monitored base by base through trityl analysis to			trityl analysis to ensure	
	appro	priate coupling effici	ency. The oligo is subsequently pu	rified by affinity-solid	
	phase	e extraction. The anno	ealed RNA duplex is further analyz	ed by mass	
	spect	rometry to verify the	exact composition of the duplex.	Each lot is compared to	
	the p	revious lot by mass s	pectrometry to ensure maximum l	ot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	mous	mouse FOXJ1 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can			
	be tra	ansfected individually	or pooled together to achieve kno	ockdown of the target	
	gene,	gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	FOXJ	I1 siRNA (Mouse) - A	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

FOXJ1 siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
FOXJ1 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
_		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μl
		100 nM	5 μl	2 μl
12-well	1 ml	50 nM	2.5 μl	2 μl
		10 nM	0.5 μl	2 μΙ
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 μΙ
		10 nM	1 μl	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com