

Product Data Sheet

FOXF1 siRNA (Mouse)

| Catalog # | Source | Reactivity | Applications | | |
|---|------------|--|--------------------------------------|--------------------------|--|
| CRM1818 | Synthetic | Μ | RNAi | | |
| Description | siRNA | to inhibit FOXF1 exp | ression using RNA interference | | |
| Specificity | FOXF1 | FOXF1 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed to | | | |
| | knock | down gene expression | on. | | |
| Form | Lyoph | ilized powder | | | |
| Gene Symbol | FOXF1 | FOXF1 | | | |
| Alternative Na | ames FKHL5 | FKHL5; FOXF1A; FREAC1; HFH8; Forkhead box protein F1; Forkhead-related protein | | | |
| | FKHL5 | 5; Forkhead-related tr | anscription factor 1; FREAC-1; Hep | patocyte nuclear factor | |
| | 3 fork | head homolog 8; HFF | 1-8 | | |
| Entrez Gene | 15227 | 7 (Mouse) | | | |
| SwissProt | Q610 | Q61080 (Mouse) | | | |
| Purity | > 97% | > 97% | | | |
| Quality Control Oligonucleotide synthesis is monitored base by base through trityl analysis | | | trityl analysis to ensure | | |
| | appro | priate coupling efficie | ency. The oligo is subsequently pu | rified by affinity-solid | |
| | phase | extraction. The anne | aled RNA duplex is further analyze | ed by mass | |
| | spect | rometry to verify the | exact composition of the duplex. I | Each lot is compared to | |
| | the pr | revious lot by mass sp | ectrometry to ensure maximum lo | ot-to-lot consistency. | |
| Components | We of | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of | | | |
| | mous | e FOXF1 gene. Each v | ial contains 5 nmol of lyophilized s | iRNA. The duplexes can | |
| | be tra | insfected individually | or pooled together to achieve kno | ockdown of the target | |
| | gene, | gene, which is most commonly assessed by qPCR or western blot. | | | |
| | Com | ponent | 15 nmol | 30 nmol | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| FOXF1 siRNA (Mouse) - A | 5 nmol x 1 | 5 nmol x 2 |
|-------------------------|--------------|--------------|
| FOXF1 siRNA (Mouse) - B | 5 nmol x 1 | 5 nmol x 2 |
| FOXF1 siRNA (Mouse) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μΙ |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| | | 10 nM | 0.25 μl | 1 µl |
| | | 100 nM | 5 μl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| | | 100 nM | 10 µl | 5 µl |
| 6-well | 2 ml | 50 nM | 5 μl | 5 μl |
| | | 10 nM | 1 μl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



For research purposes only, not for human use

Product Data Sheet

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com