

Product Data Sheet

KCNMB1 siRNA (Mouse)

| Catalog # | Source | Reactivity | Applications | | | |
|----------------|--|--|---|--|--|--|
| CRM2191 | Synthetic | Μ | RNAi | | | |
| Description | siRNA | to inhibit KCNMB1 e | xpression using RNA interference | | | |
| Specificity | KCNIV | 1B1 siRNA (Mouse) is | a target-specific 19-23 nt siRNA oligo duplexes designed | | | |
| | to kno | ock down gene expre | ssion. | | | |
| Form | Lyoph | ilized powder | | | | |
| Gene Symbol | KCNIV | KCNMB1 | | | | |
| Alternative Na | mes Calciu | Calcium-activated potassium channel subunit beta-1; BK channel subunit beta-1; | | | | |
| | BKbet | a; BKbeta1; Calcium- | activated potassium channel. subfamily M subunit beta-1; | | | |
| | Calciu | im-activated potassiu | m channel subunit beta; Charybdotoxin receptor subunit | | | |
| | beta-: | 1; K(VCA | | | | |
| Entrez Gene | 16533 | 16533 (Mouse) | | | | |
| SwissProt | Q8CA | Q8CAE3 (Mouse) | | | | |
| Purity | > 97% | ,) | | | | |
| Quality Contro | l Oligor | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure | | | | |
| | appro | priate coupling effici | ency. The oligo is subsequently purified by affinity-solid | | | |
| | phase | e extraction. The anne | ealed RNA duplex is further analyzed by mass | | | |
| | spect | rometry to verify the | exact composition of the duplex. Each lot is compared to | | | |
| | the pi | revious lot by mass sp | pectrometry to ensure maximum lot-to-lot consistency. | | | |
| Components | We of | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of | | | | |
| | mous | e KCNMB1 gene. Eac | n vial contains 5 nmol of lyophilized siRNA. The duplexes | | | |
| | can b | e transfected individu | ally or pooled together to achieve knockdown of the | | | |
| | target | t gene, which is most | commonly assessed by qPCR or western blot. | | | |
| Components | spect the pr We of mous can be target | rometry to verify the revious lot by mass s ffers pre-designed se e KCNMB1 gene. Eac e transfected individu t gene, which is most | exact composition of the duplex. Each lot is compared to bectrometry to ensure maximum lot-to-lot consistency. Its of 3 different target-specific siRNA oligo duplexes of in vial contains 5 nmol of lyophilized siRNA. The duplexes wally or pooled together to achieve knockdown of the | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| Component | 15 nmol | 30 nmol |
|--------------------------|--------------|--------------|
| KCNMB1 siRNA (Mouse) - A | 5 nmol x 1 | 5 nmol x 2 |
| KCNMB1 siRNA (Mouse) - B | 5 nmol x 1 | 5 nmol x 2 |
| KCNMB1 siRNA (Mouse) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 μl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 µl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| | | 10 nM | 0.25 μl | 1 μΙ |
| | | 100 nM | 5 µl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| 6-well | 2 ml | 100 nM | 10 µl | 5 µl |
| | | 50 nM | 5 µl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com