

Product Data Sheet

SLC22A18 siRNA (Mouse)

| Catalog # | Source | Reactivity | Applications | | | |
|--------------------------------|-----------|---|---|--|--|--|
| CRM2915 | Synthetic | M | RNAi | | | |
| Description | siRN | A to inhibit SLC22A18 | expression using RNA interference | | | |
| Specificity | SLC2 | 2A18 siRNA (Mouse) | is a target-specific 19-23 nt siRNA oligo duplexes designed | | | |
| | to kn | lock down gene expre | ession. | | | |
| Form | Lyop | hilized powder | | | | |
| Gene Symbol | SLC2 | SLC22A18 | | | | |
| Alternative Na | ames IMPT | IMPT1; ITM; ORCTL2; TSSC5; Solute carrier family 22 member 18; Imprinted | | | | |
| | multi | i-membrane-spannin | g polyspecific transporter-related protein 1; | | | |
| | Mult | i-membrane-spannin | g polyspecific transporter; Organic cation transporter-like | | | |
| | prote | ein 2; ORCTL-2 | | | | |
| Entrez Gene | 1840 | 18400 (Mouse) | | | | |
| SwissProt | Q78k | Q78KK3 (Mouse) | | | | |
| Purity | > 97% | % | | | | |
| Quality Control | ol Oligo | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure | | | | |
| | appro | opriate coupling effici | iency. The oligo is subsequently purified by affinity-solid | | | |
| | phas | e extraction. The ann | ealed RNA duplex is further analyzed by mass | | | |
| | spect | trometry to verify the | exact composition of the duplex. Each lot is compared to | | | |
| | the p | previous lot by mass s | pectrometry to ensure maximum lot-to-lot consistency. | | | |
| Components | We o | offers pre-designed se | ts of 3 different target-specific siRNA oligo duplexes of | | | |
| | mous | se SLC22A18 gene. Ea | ch vial contains 5 nmol of lyophilized siRNA. The duplexes | | | |
| | can b | e transfected individ | ually or pooled together to achieve knockdown of the | | | |
| | targe | t gene, which is most | commonly assessed by qPCR or western blot. | | | |
| A secold sectors in the second | | | abiata abamiatan JE Janan mafluana ana EC Elavianta ana ta | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| Component | 15 nmol | 30 nmol |
|----------------------------|--------------|--------------|
| SLC22A18 siRNA (Mouse) - A | 5 nmol x 1 | 5 nmol x 2 |
| SLC22A18 siRNA (Mouse) - B | 5 nmol x 1 | 5 nmol x 2 |
| SLC22A18 siRNA (Mouse) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 μl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 µl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μl |
| | | 10 nM | 0.25 μl | 1 μl |
| | | 100 nM | 5 µl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| 6-well | 2 ml | 100 nM | 10 µl | 5 µl |
| | | 50 nM | 5 µl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com