

Product Data Sheet

ENPP1 siRNA (Mouse)

| Catalog #SourceReactivityApplicationsCRM3013SyntheticMRNAiDescriptionsiRNA to inhibit ENPP1 expression using RNA interferenceSpecificityENPP1 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed knock down gene expression.FormLyophilized powderGene SymbolENPP1Alternative NamesNPPS; PC1; PDNP1; Ectonucleotide pyrophosphatase/phosphodiesterase family member 1; E-NPP 1; Lymphocyte antigen 41; Ly-41; Phosphodiesterase l/nucleot pyrophosphatase 1; Plasma-cell membrane glycoprotein PC-1Entrez Gene18605 (Mouse)SwissProtP06802 (Mouse)Purity> 97%Quality ControlOligonucleotide synthesis is monitored base by base through trityl analysis to en appropriate coupling efficiency. The oligo is subsequently purified by affinity-soli phase extraction. The annealed RNA duplex is further analyzed by mass | | | |
|--|--|--|--|
| DescriptionsiRNA to inhibit ENPP1 expression using RNA interferenceSpecificityENPP1 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed knock down gene expression.FormLyophilized powderGene SymbolENPP1Alternative NamesNPPS; PC1; PDNP1; Ectonucleotide pyrophosphatase/phosphodiesterase family member 1; E-NPP 1; Lymphocyte antigen 41; Ly-41; Phosphodiesterase l/nucleot pyrophosphatase 1; Plasma-cell membrane glycoprotein PC-1Entrez Gene18605 (Mouse)SwissProtP06802 (Mouse)Purity> 97%Quality ControlOligonucleotide synthesis is monitored base by base through trityl analysis to en appropriate coupling efficiency. The oligo is subsequently purified by affinity-soli | | | |
| SpecificityENPP1 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed knock down gene expression.FormLyophilized powderGene SymbolENPP1Alternative NamesNPPS; PC1; PDNP1; Ectonucleotide pyrophosphatase/phosphodiesterase family member 1; E-NPP 1; Lymphocyte antigen 41; Ly-41; Phosphodiesterase l/nucleot pyrophosphatase 1; Plasma-cell membrane glycoprotein PC-1Entrez Gene18605 (Mouse)SwissProtP06802 (Mouse)Purity> 97%Quality ControlOligonucleotide synthesis is monitored base by base through trityl analysis to en appropriate coupling efficiency. The oligo is subsequently purified by affinity-soli | | | |
| knock down gene expression.FormLyophilized powderGene SymbolENPP1Alternative NamesNPPS; PC1; PDNP1; Ectonucleotide pyrophosphatase/phosphodiesterase family member 1; E-NPP 1; Lymphocyte antigen 41; Ly-41; Phosphodiesterase l/nucleot pyrophosphatase 1; Plasma-cell membrane glycoprotein PC-1Entrez Gene18605 (Mouse)SwissProtP06802 (Mouse)Purity> 97%Quality ControlOligonucleotide synthesis is monitored base by base through trityl analysis to en appropriate coupling efficiency. The oligo is subsequently purified by affinity-soli | | | |
| FormLyophilized powderGene SymbolENPP1Alternative NamesNPPS; PC1; PDNP1; Ectonucleotide pyrophosphatase/phosphodiesterase family member 1; E-NPP 1; Lymphocyte antigen 41; Ly-41; Phosphodiesterase l/nucleot pyrophosphatase 1; Plasma-cell membrane glycoprotein PC-1Entrez Gene18605 (Mouse)SwissProtP06802 (Mouse)Purity> 97%Quality ControlOligonucleotide synthesis is monitored base by base through trityl analysis to en appropriate coupling efficiency. The oligo is subsequently purified by affinity-soli | | | |
| Gene SymbolENPP1Alternative NamesNPPS; PC1; PDNP1; Ectonucleotide pyrophosphatase/phosphodiesterase family member 1; E-NPP 1; Lymphocyte antigen 41; Ly-41; Phosphodiesterase l/nucleot pyrophosphatase 1; Plasma-cell membrane glycoprotein PC-1Entrez Gene18605 (Mouse)SwissProtP06802 (Mouse)Purity> 97%Quality ControlOligonucleotide synthesis is monitored base by base through trityl analysis to en appropriate coupling efficiency. The oligo is subsequently purified by affinity-soli | | | |
| Alternative NamesNPPS; PC1; PDNP1; Ectonucleotide pyrophosphatase/phosphodiesterase family member 1; E-NPP 1; Lymphocyte antigen 41; Ly-41; Phosphodiesterase l/nucleot pyrophosphatase 1; Plasma-cell membrane glycoprotein PC-1Entrez Gene18605 (Mouse)SwissProtP06802 (Mouse)Purity> 97%Quality ControlOligonucleotide synthesis is monitored base by base through trityl analysis to en appropriate coupling efficiency. The oligo is subsequently purified by affinity-soli | | | |
| member 1; E-NPP 1; Lymphocyte antigen 41; Ly-41; Phosphodiesterase I/nucleot pyrophosphatase 1; Plasma-cell membrane glycoprotein PC-1Entrez Gene18605 (Mouse)SwissProtP06802 (Mouse)Purity> 97%Quality ControlOligonucleotide synthesis is monitored base by base through trityl analysis to en appropriate coupling efficiency. The oligo is subsequently purified by affinity-soli | | | |
| pyrophosphatase 1; Plasma-cell membrane glycoprotein PC-1Entrez Gene18605 (Mouse)SwissProtP06802 (Mouse)Purity> 97%Quality ControlOligonucleotide synthesis is monitored base by base through trityl analysis to en appropriate coupling efficiency. The oligo is subsequently purified by affinity-soli | | | |
| Entrez Gene18605 (Mouse)SwissProtP06802 (Mouse)Purity> 97%Quality ControlOligonucleotide synthesis is monitored base by base through trityl analysis to en appropriate coupling efficiency. The oligo is subsequently purified by affinity-soli | | | |
| SwissProt P06802 (Mouse) Purity > 97% Quality Control Oligonucleotide synthesis is monitored base by base through trityl analysis to en appropriate coupling efficiency. The oligo is subsequently purified by affinity-soli | | | |
| Purity > 97% Quality Control Oligonucleotide synthesis is monitored base by base through trityl analysis to en appropriate coupling efficiency. The oligo is subsequently purified by affinity-soli | | | |
| Quality ControlOligonucleotide synthesis is monitored base by base through trityl analysis to en appropriate coupling efficiency. The oligo is subsequently purified by affinity-soli | | | |
| appropriate coupling efficiency. The oligo is subsequently purified by affinity-soli | | | |
| | | | |
| phase extraction. The annealed RNA duplex is further analyzed by mass | | | |
| | | | |
| spectrometry to verify the exact composition of the duplex. Each lot is compared the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency | | | |
| | | | |
| mouse ENPP1 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes | | | |
| be transfected individually or pooled together to achieve knockdown of the targe | | | |
| gene, which is most commonly assessed by qPCR or western blot. | | | |
| Component 15 nmol 30 nmol | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| ENPP1 siRNA (Mouse) - A | 5 nmol x 1 | 5 nmol x 2 |
|-------------------------|--------------|--------------|
| ENPP1 siRNA (Mouse) - B | 5 nmol x 1 | 5 nmol x 2 |
| ENPP1 siRNA (Mouse) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μΙ |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| | | 10 nM | 0.25 μl | 1 μΙ |
| | | 100 nM | 5 μl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| | | 100 nM | 10 µl | 5 μΙ |
| 6-well | 2 ml | 50 nM | 5 μl | 5 μΙ |
| | | 10 nM | 1 μΙ | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



For research purposes only, not for human use

Product Data Sheet

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com