

Product Data Sheet

POU3F2 siRNA (Mouse)

| Catalog # | Source | Reactivity | Applications | | | |
|---------------|-----------|---|---|--|--|--|
| CRM3168 | Synthetic | М | RNAi | | | |
| Description | siRN | A to inhibit POU3F2 e | xpression using RNA interference | | | |
| Specificity | POU | POU3F2 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed to | | | | |
| | knoc | k down gene express | on. | | | |
| Form | Lyop | hilized powder | | | | |
| Gene Symbol | POU | POU3F2 | | | | |
| Alternative N | ames BRN- | BRN-2; BRN2; OTF7; POU domain. class 3. transcription factor 2; Brain-specific | | | | |
| | hom | eobox/POU domain p | rotein 2; Brain-2; Brn-2; Nervous system-specific | | | |
| | octai | mer-binding transcrip | tion factor N-Oct-3; Octamer-binding protein 7; Oct-7; | | | |
| | Octa | mer-binding transcrip | | | | |
| Entrez Gene | 1899 | 18992 (Mouse) | | | | |
| SwissProt | P313 | P31360 (Mouse) | | | | |
| Purity | > 979 | % | | | | |
| Quality Contr | ol Oligo | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure | | | | |
| | appr | opriate coupling effic | iency. The oligo is subsequently purified by affinity-solid | | | |
| | phas | e extraction. The ann | ealed RNA duplex is further analyzed by mass | | | |
| | spec | trometry to verify the | exact composition of the duplex. Each lot is compared to | | | |
| | the p | previous lot by mass s | pectrometry to ensure maximum lot-to-lot consistency. | | | |
| Components | We c | offers pre-designed se | ts of 3 different target-specific siRNA oligo duplexes of | | | |
| | mou | se POU3F2 gene. Eacl | n vial contains 5 nmol of lyophilized siRNA. The duplexes | | | |
| | can b | pe transfected individ | ually or pooled together to achieve knockdown of the | | | |
| | targe | et gene, which is most | commonly assessed by qPCR or western blot. | | | |
| | | | | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| Component | 15 nmol | 30 nmol |
|--------------------------|--------------|--------------|
| POU3F2 siRNA (Mouse) - A | 5 nmol x 1 | 5 nmol x 2 |
| POU3F2 siRNA (Mouse) - B | 5 nmol x 1 | 5 nmol x 2 |
| POU3F2 siRNA (Mouse) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 µM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 μl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 µl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μl |
| | | 10 nM | 0.25 μl | 1 µl |
| | | 100 nM | 5 µl | 2 µl |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| 6-well | 2 ml | 100 nM | 10 µl | 5 µl |
| | | 50 nM | 5 µl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com