

Product Data Sheet

POU4F1 siRNA (Mouse)

| Catalog # | Source | Reactivity | Applications | | |
|--|------------|--|------------------------------------|---------------------------|--|
| CRM3171 | Synthetic | Μ | RNAi | | |
| Description | siRNA | to inhibit POU4F1 ex | pression using RNA interference | | |
| Specificity | POU4 | F1 siRNA (Mouse) is a | a target-specific 19-23 nt siRNA o | ligo duplexes designed to | |
| | knock | down gene expression | on. | | |
| Form | Lyoph | ilized powder | | | |
| Gene Symbol | POU4 | POU4F1 | | | |
| Alternative N | ames BRN-3 | BRN-3; BRN3; BRN3A; POU domain. class 4. transcription factor 1; Brain-specific | | | |
| | home | obox/POU domain p | rotein 3A; Brain-3A; Brn-3A; Brn-3 | 3.0 | |
| Entrez Gene | 18996 | (Mouse) | | | |
| SwissProt P17208 (Mouse) | | | | | |
| Purity > 97% | | | | | |
| Quality Control Oligonucleotide synthesis is monitored base by base through trit | | | h trityl analysis to ensure | | |
| | appro | priate coupling effici | ency. The oligo is subsequently p | urified by affinity-solid | |
| | phase | extraction. The anne | ealed RNA duplex is further analy: | zed by mass | |
| | spectr | ometry to verify the | exact composition of the duplex. | Each lot is compared to | |
| | the pr | evious lot by mass sp | pectrometry to ensure maximum | lot-to-lot consistency. | |
| Components | We of | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of | | | |
| | mouse | e POU4F1 gene. Each | vial contains 5 nmol of lyophilize | d siRNA. The duplexes | |
| | can be | e transfected individu | ally or pooled together to achiev | e knockdown of the | |
| | target | target gene, which is most commonly assessed by qPCR or western blot. | | | |
| | Com | ponent | 15 nmol | 30 nmol | |
| | POU | 4F1 siRNA (Mouse) - | A 5 nmol x 1 | 5 nmol x 2 | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| POU4F1 siRNA (Mouse) - B | 5 nmol x 1 | 5 nmol x 2 |
|--------------------------|--------------|--------------|
| POU4F1 siRNA (Mouse) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| | | 10 nM | 0.25 μl | 1 µl |
| | | 100 nM | 5 μl | 2 µl |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| | | 100 nM | 10 µl | 5 µl |
| 6-well | 2 ml | 50 nM | 5 μl | 5 μΙ |
| | | 10 nM | 1 µl | 5 μΙ |

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |