

# **Product Data Sheet**

### PRL8A6 siRNA (Mouse)

Catalog #	Source	Reactivity	Applications			
CRM3232	Synthetic	Μ	RNAi			
Description	siRNA	to inhibit PRL8A6 ex	pression using RNA interference			
Specificity	PRL8A	PRL8A6 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed to				
	knock	down gene expressi	on.			
Form	Lyoph	Lyophilized powder				
Gene Symbol	PRL8A	PRL8A6				
Alternative N	ames PRLPC	PRLPC; PRLPC1; Prolactin-8A6; Placental prolactin-like protein C1; PLP-C1; PRL-like				
	protei	in C1; Prolactin-like p	rotein C-alpha; PLP C-alpha; Pro	lactin-like protein C		
Entrez Gene	19112	19112 (Mouse)				
SwissProt	Q9DA	Y2 (Mouse)				
Purity > 9		> 97%				
Quality Contr	Quality Control Oligonucleotide synthesis is monitored base by base through trityl analysis to			gh trityl analysis to ensure		
	appro	appropriate coupling efficiency. The oligo is subsequently purified by affinity-se				
	phase	extraction. The ann	ealed RNA duplex is further analy	yzed by mass		
	spectr	rometry to verify the	exact composition of the duplex	. Each lot is compared to		
	the pr	revious lot by mass s	pectrometry to ensure maximum	n lot-to-lot consistency.		
Components We offers pre-designed sets of 3 different			ts of 3 different target-specific si	ent target-specific siRNA oligo duplexes of		
	mouse	e PRL8A6 gene. Each	vial contains 5 nmol of lyophilize	ed siRNA. The duplexes		
	can be	e transfected individ	ually or pooled together to achie	ve knockdown of the		
	target	target gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent	15 nmol	30 nmol		
	PRL8	A6 siRNA (Mouse)	A 5 nmol x 1	5 nmol x 2		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



# **Product Data Sheet**

PRL8A6 siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
PRL8A6 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 μl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 μΙ
		10 nM	1 μl	5 μΙ

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com