

Product Data Sheet

UPF1 siRNA (Mouse)

| Catalog # | Source | Reactivity | Applications | | |
|----------------|------------|---|-----------------------------------|--------------------------|--|
| CRM3457 | Synthetic | М | RNAi | | |
| Description | siRNA | to inhibit UPF1 expres | ssion using RNA interference | | |
| Specificity | UPF1 | siRNA (Mouse) is a tar | get-specific 19-23 nt siRNA oligo | duplexes designed to | |
| | knock | down gene expression | ۱. | | |
| Form | Lyoph | ilized powder | | | |
| Gene Symbol | UPF1 | UPF1 | | | |
| Alternative Na | ames RENT1 | RENT1; Regulator of nonsense transcripts 1; ATP-dependent helicase RENT1; | | | |
| | Nonse | ense mRNA reducing fa | actor 1; NORF1; Up-frameshift su | ppressor 1 homolog; | |
| | mUpf: | 1 | | | |
| Entrez Gene | 19704 | (Mouse) | | | |
| SwissProt | Q9EPU | Q9EPU0 (Mouse) | | | |
| Purity | > 97% | > 97% | | | |
| Quality Contro | ol Oligor | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure | | | |
| | appro | priate coupling efficier | ncy. The oligo is subsequently pu | rified by affinity-solid | |
| | phase | extraction. The annea | led RNA duplex is further analyz | ed by mass | |
| | spectr | ometry to verify the e | xact composition of the duplex. | Each lot is compared to | |
| | the pr | evious lot by mass spe | ectrometry to ensure maximum l | ot-to-lot consistency. | |
| Components | We of | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of | | | |
| | mouse | e UPF1 gene. Each vial | contains 5 nmol of lyophilized si | iRNA. The duplexes can | |
| | be tra | nsfected individually o | r pooled together to achieve kno | ockdown of the target | |
| | gene, | gene, which is most commonly assessed by qPCR or western blot. | | | |
| | Com | ponent | 15 nmol | 30 nmol | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| UPF1 siRNA (Mouse) - A | 5 nmol x 1 | 5 nmol x 2 |
|------------------------|--------------|--------------|
| UPF1 siRNA (Mouse) - B | 5 nmol x 1 | 5 nmol x 2 |
| UPF1 siRNA (Mouse) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 µl of DEPC water to get a final concentration of 20 µM.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μl |
| | | 10 nM | 0.25 μl | 1 μl |
| | | 100 nM | 5 µl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| | | 100 nM | 10 µl | 5 μΙ |
| 6-well | 2 ml | 50 nM | 5 µl | 5 μl |
| | | 10 nM | 1 µl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



For research purposes only, not for human use

Product Data Sheet

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com