

# **Product Data Sheet**

## **RFX2 siRNA (Mouse)**

Catalog #	Source	Reactivity	Applications		
CRM3469	Synthetic	М	RNAi		
<b>Description</b> siRNA to inhibit RFX2 expression using RNA inte			ssion using RNA interference		
Specificity	RFX2	RFX2 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed to			
	knock	knock down gene expression.			
Form	Lyoph	Lyophilized powder			
Gene Symbol	RFX2	RFX2			
Alternative Names DNA-binding protein RFX2; Regulatory factor X 2					
Entrez Gene 19725 (Mouse)					
SwissProt	P4837	P48379 (Mouse)			
Purity > 97%					
Quality Control Oligonucleotide synthesis is monitored base by base through trityl analysis			n trityl analysis to ensure		
appropriate coupling efficiency. The oligo is subsequently purified by affin			rified by affinity-solid		
phase extraction. The annealed RNA duplex is further analyzed by m			ed by mass		
	spect	spectrometry to verify the exact composition of the duplex. Each lot is compared to			
	the pi	revious lot by mass sp	ectrometry to ensure maximum l	ot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	mous	e RFX2 gene. Each via	l contains 5 nmol of lyophilized si	RNA. The duplexes can	
	be tra	insfected individually	or pooled together to achieve kno	ockdown of the target	
gene, which is most commonly assessed by qPCR or western blot.			ı blot.		
		ponent	15 nmol	30 nmol	
	RFX2	2 siRNA (Mouse) - A	5 nmol x 1	5 nmol x 2	

 RFX2 siRNA (Mouse) - B
 5 nmol x 1
 5 nmol x 2

 Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC 

 Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility

 Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

 Species reactivity key: H- Human, Ma Mouse, R- Bat, B- Boying, C- Chicken, D- Dog, G- Goat, Mk, Monkey, P- Pig, Bh 

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

L	DEPC Water	T 1111 X T	
F	NERC Water	1 ml x 1	1 ml x 2
٩	Negative Control	2.5 nmol x 1	2.5 nmol x 2
F	RFX2 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 µl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 μl

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com