

Product Data Sheet

SEMA4B siRNA (Mouse)

Catalog #	Source	Reactivity	Арр	lications	
CRM3665	Synthetic	М	RNA	i	
Description	siRNA	to inhibit SEMA4B e	xpression using RNA i	nterference	
Specificity	SEMA	4B siRNA (Mouse) is	a target-specific 19-23	3 nt siRNA oligo duplexes designed	
	to kno	ock down gene expre	ssion.		
Form	Lyoph	nilized powder			
Gene Symbol	SEMA	SEMA4B			
Alternative N	ames KIAA1	KIAA1745; SEMAC; SEMC; Semaphorin-4B; Semaphorin-C; Sema C			
Entrez Gene	20352	2 (Mouse)			
SwissProt	Q621	79 (Mouse)			
Purity	> 97%	6			
Quality Contr	ol Oligo	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure			
	appro	opriate coupling effici	ency. The oligo is subs	sequently purified by affinity-solid	
	phase	e extraction. The ann	ealed RNA duplex is fu	irther analyzed by mass	
	spect	rometry to verify the	exact composition of	the duplex. Each lot is compared to	
	the p	revious lot by mass s	pectrometry to ensure	e maximum lot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	mous	e SEMA4B gene. Eac	n vial contains 5 nmol	of lyophilized siRNA. The duplexes	
	can b	can be transfected individually or pooled together to achieve knockdown of the			
	targe	target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nn	nol 30 nmol	
	SEM	A4B siRNA (Mouse) -	A 5 nmc	bl x 1 5 nmol x 2	
	SEM	A4B siRNA (Mouse) -	B 5 nmo	bl x 1 5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

DEPC	CWater	1 ml x 1	1 ml x 2
Nega	tive Control	2.5 nmol x 1	2.5 nmol x 2
SEM	A4B siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com