

Product Data Sheet

ST6GALNAC1 siRNA (Mouse)

Catalog #	Source	Reactivity	Applicatio	ns	
CRM3720	Synthetic	М	RNAi		
Description	siRNA	to inhibit ST6GALN	AC1 expression using RNA int	terference	
Specificity	ST6G	ALNAC1 siRNA (Mou	se) is a target-specific 19-23	nt siRNA oligo duplexes	
	desig	ned to knock down <code>g</code>	ene expression.		
Form	Lyoph	Lyophilized powder			
Gene Symbol	ST6G	ST6GALNAC1			
Alternative N	ames SIAT7	SIAT7A; Alpha-N-acetylgalactosaminide alpha-2.6-sialyltransferase 1; GalNAc			
	alpha	-2.6-sialyltransferase	e I; ST6GalNAc I; ST6GalNAcI;	; Sialyltransferase 7A; SIAT7-A	
Entrez Gene	2044	20445 (Mouse)			
SwissProt C		Q9QZ39 (Mouse)			
Purity > 97%		6			
Quality Control Oligonucleotide synthesis is monitored base by base through trityl analysis to			hrough trityl analysis to ensure		
	appro	opriate coupling effic	iency. The oligo is subsequer	ntly purified by affinity-solid	
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass			
	spect	spectrometry to verify the exact composition of the duplex. Each lot is compared to			
	the p	the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.			
Components	We o	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	mous	mouse ST6GALNAC1 gene. Each vial contains 5 nmol of lyophilized siRNA. The			
	duple	duplexes can be transfected individually or pooled together to achieve knockdown			
	of the	of the target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	ST60	GALNAC1 siRNA (Mo	use) - A 5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com

Product Data Sheet

ST6GA	LNAC1 siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
ST6GA	LNAC1 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negati	ve Control	2.5 nmol x 1	2.5 nmol x 2
DEPC	Nater	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μΙ
24-well	500 μl	50 nM	1.25 μl	1 μl
_		10 nM	0.25 μl	1 μΙ
		100 nM	5 µl	2 μΙ
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 µl	5 µl
		10 nM	1 μΙ	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com