

Product Data Sheet

WNT5B siRNA (Mouse)

Catalog #	Source	Reactivity	Applications		
CRM4435	Synthetic	М	RNAi		
Description	siRNA	to inhibit WNT5B ex	pression using RNA interferenc	ie	
Specificity	WNT	5B siRNA (Mouse) is a	a target-specific 19-23 nt siRNA	oligo duplexes designed to	
	knock	k down gene expressi	on.		
Form	Lyoph	nilized powder			
Gene Symbol	WNT	5B			
Alternative N	ames WNT-	WNT-5B; Protein Wnt-5b			
Entrez Gene	22419	9 (Mouse)			
SwissProt	P2272	26 (Mouse)			
Purity	> 97%	> 97%			
Quality Contr	ol Oligo	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure			
	appro	opriate coupling effici	ency. The oligo is subsequently	purified by affinity-solid	
	phase	e extraction. The anno	ealed RNA duplex is further ana	alyzed by mass	
	spect	rometry to verify the	exact composition of the duple	ex. Each lot is compared to	
	the p	revious lot by mass s	pectrometry to ensure maximu	m lot-to-lot consistency.	
Components	We of	ffers pre-designed se	ts of 3 different target-specific	siRNA oligo duplexes of	
	mous	e WNT5B gene. Each	vial contains 5 nmol of lyophili	zed siRNA. The duplexes	
	can b	e transfected individu	ually or pooled together to achi	eve knockdown of the	
	target gene, which is most commonly assessed by qPCR or western blot.			r western blot.	
	Com	ponent	15 nmol	30 nmol	
	WN	Г5B siRNA (Mouse) - л	A 5 nmol x 1	5 nmol x 2	
	WN	T5B siRNA (Mouse) - ∣	B 5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

WNT5B siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
of medium	of siRNA		2000
	100 nM	0.5 μl	0.25 μl
100 µl	50 nM	0.25 μl	0.25 μl
	10 nM	0.05 μl	0.25 μl
	100 nM	2.5 μl	1 µl
500 μl	50 nM	1.25 μl	1 µl
	10 nM	0.25 μl	1 µl
	100 nM	5 µl	2 µl
1 ml	50 nM	2.5 μl	2 µl
	10 nM	0.5 μl	2 µl
	100 nM	10 µl	5 µl
2 ml	50 nM	5 μl	5 µl
	10 nM	1 µl	5 µl
	of medium 100 μl 500 μl 1 ml	of medium of siRNA 100 nM 100 nM 100 nM 10 nM 50 nM 10 nM 500 μl 50 nM 100 nM 10 nM 500 μl 50 nM 10 nM 10 nM 10 nM 10 nM 10 nM 10 nM 10 nM 10 nM 1 nn 50 nM 10 nM 10 nM 10 nM 50 nM	of mediumof siRNA100 nM0.5 μl100 μl50 nM0.25 μl10 nM0.05 μl500 μl10 nM2.5 μl500 μl50 nM1.25 μl10 nM0.25 μl10 nM0.25 μl10 nM0.25 μl10 nM5 μl100 nM5 μl100 nM10 μl100 nM10 μl100 nM5.0 μl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com