

Product Data Sheet

ZSCAN12 siRNA (Mouse)

Catalog #	Source	Reactivity		Applications		
CRM4501	Synthetic	Μ		RNAi		
Description siRNA to inhibit ZSCA			expression using F	RNA interference		
Specificity	ZSCAI	ZSCAN12 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed				
	to kno	ock down gene expre	ssion.			
Form	Lyoph	Lyophilized powder				
Gene Symbol	ZSCAI	ZSCAN12				
Alternative N	ames ZFP96	ZFP96; ZNF96; Zinc finger and SCAN domain-containing protein 12; Zinc finger				
	prote	in 96; Zfp-96				
Entrez Gene	22758	22758 (Mouse)				
SwissProt	Q9Z1	Q9Z1D7 (Mouse)				
Purity	> 97%	> 97%				
Quality Contr	ol Oligoi	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure				
	appro	appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid				
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass				
	spect	spectrometry to verify the exact composition of the duplex. Each lot is compared to				
	the p	the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.				
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of				
	mous	e ZSCAN12 gene. Eac	h vial contains 5 r	nmol of lyophilized siRNA. The duplex	es	
	can b	e transfected individu	ally or pooled to	gether to achieve knockdown of the		
	target	target gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent	1	5 nmol 30 nmol		
	ZSCA	N12 siRNA (Mouse)	- A 5	nmol x 1 5 nmol x 2		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

ZSCAN12 siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
ZSCAN12 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
_		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μl
		10 nM	0.25 μl	1 μl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 μl
_		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 μΙ
		10 nM	1 μΙ	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com