

# **Product Data Sheet**

## SLC27A5 siRNA (Mouse)

| Catalog #            | Source    | Reactivity   | Applications  |  |  |
|----------------------|-----------|--|---|--|--|
| CRM4760              | Synthetic | Μ  | RNAi  |  |  |
| Description          | siRNA     | ι to inhibit SLC27A5 ε   | expression using RNA interference   |  |  |
| Specificity          | SLC27     | 7A5 siRNA (Mouse) is   | a target-specific 19-23 nt siRNA oligo duplexes designed  |  |  |
|                      | to kn     | ock down gene expre  | ession.   |  |  |
| Form                 | Lyopł     | nilized powder   |   |  |  |
| Gene Symbol          | SLC2      | SLC27A5  |   |  |  |
| Alternative Na       | ames ACSB | ACSB; ACSVL6; FATP5; VLACSR; Bile acyl-CoA synthetase; BACS; Bile acid-CoA ligase; |   |  |  |
|                      | BA-Co     | oA ligase; BAL; Cholat   | eCoA ligase; Fatty acid transport protein 5; FATP-5;  |  |  |
|                      | Solut     | e carrier family 27 m  | ember 5; Very long-chain acyl-CoA synthetase-related  |  |  |
|                      | prote     | ein; VLA   |   |  |  |
| Entrez Gene          | 2645      | 26459 (Mouse)  |   |  |  |
| SwissProt            | Q4LD      | Q4LDG0 (Mouse)   |   |  |  |
| Purity               | > 97%     | 6  |   |  |  |
| Quality Contro       | ol Oligo  | nucleotide synthesis   | is monitored base by base through trityl analysis to ensure   |  |  |
|                      | appro     | opriate coupling effici  | ency. The oligo is subsequently purified by affinity-solid  |  |  |
|                      | phase     | e extraction. The ann  | ealed RNA duplex is further analyzed by mass  |  |  |
|                      | spect     | rometry to verify the  | exact composition of the duplex. Each lot is compared to  |  |  |
|                      | the p     | revious lot by mass s  | pectrometry to ensure maximum lot-to-lot consistency.   |  |  |
| Components           | We o      | ffers pre-designed se  | ts of 3 different target-specific siRNA oligo duplexes of   |  |  |
|                      | mous      | se SLC27A5 gene. Eac   | h vial contains 5 nmol of lyophilized siRNA. The duplexes   |  |  |
|                      | can b     | e transfected individ  | ually or pooled together to achieve knockdown of the  |  |  |
|                      | targe     | t gene, which is most  | commonly assessed by qPCR or western blot.  |  |  |
| A surface the second |           | Vesteve blet III Justeve   | a historia da la la la successiva a flucado a successiva de la successiva d |  |  |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



## **Product Data Sheet**

| Component                 | 15 nmol      | 30 nmol      |
|---------------------------|--------------|--------------|
| SLC27A5 siRNA (Mouse) - A | 5 nmol x 1   | 5 nmol x 2   |
| SLC27A5 siRNA (Mouse) - B | 5 nmol x 1   | 5 nmol x 2   |
| SLC27A5 siRNA (Mouse) - C | 5 nmol x 1   | 5 nmol x 2   |
| Negative Control          | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water                | 1 ml x 1     | 1 ml x 2     |

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

| Plate   | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
|         | of medium    | of siRNA            |               | 2000         |
|         |              | 100 nM              | 0.5 μl        | 0.25 μl      |
| 96-well | 100 μl       | 50 nM               | 0.25 μl       | 0.25 μl      |
|         |              | 10 nM               | 0.05 μl       | 0.25 μl      |
|         |              | 100 nM              | 2.5 μl        | 1 µl         |
| 24-well | 500 μl       | 50 nM               | 1.25 μl       | 1 μΙ         |
|         |              | 10 nM               | 0.25 μl       | 1 μl         |
|         |              | 100 nM              | 5 μl          | 2 μΙ         |
| 12-well | 1 ml         | 50 nM               | 2.5 μl        | 2 μΙ         |
|         |              | 10 nM               | 0.5 μl        | 2 μΙ         |
| 6-well  | 2 ml         | 100 nM              | 10 µl         | 5 µl         |
| b-Well  | 2 ml         | 50 nM               | 5 μl          | 5 μΙ         |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



## **Product Data Sheet**

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com