

# **Product Data Sheet**

### **COPS5 siRNA (Mouse)**

Catalog #	Source	Reactivity	Applications		
CRM4781	Synthetic	М	RNAi		
Description	siRNA	to inhibit COPS5 exp	ression using RNA interference		
Specificity	COPS	COPS5 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed to			
	knock	down gene expression	on.		
Form	Lyoph	ilized powder			
Gene Symbol	COPS	COPS5			
Alternative N	ames CSN5	CSN5; JAB1; KIC2; COP9 signalosome complex subunit 5; SGN5; Signalosome subunit			
	5; Jun	activation domain-b	inding protein 1; Kip1 C-terminus	-interacting protein 2	
Entrez Gene	26754	4 (Mouse)			
SwissProt C		O35864 (Mouse)			
Purity	> 97%	> 97%			
Quality Control Oligonucleotide synthesis is monitored base by base through trityl analysis t			h trityl analysis to ensure		
	appro	priate coupling effici	ency. The oligo is subsequently p	urified by affinity-solid	
	phase	e extraction. The anne	ealed RNA duplex is further analy	zed by mass	
	spect	rometry to verify the	exact composition of the duplex.	Each lot is compared to	
	the p	revious lot by mass sp	pectrometry to ensure maximum	lot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	mous	mouse COPS5 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can			
	be tra	be transfected individually or pooled together to achieve knockdown of the target			
	gene,	gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	COP	S5 siRNA (Mouse) - A	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



# **Product Data Sheet**

COPS5 siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
COPS5 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 µl
		100 nM	5 µl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 µl	5 μΙ
		10 nM	1 μΙ	5 μΙ

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com