

# **Product Data Sheet**

## FOLH1 siRNA (Mouse)

| Catalog #      | Source    | Reactivity                                                                            | Applications                                               |  |  |  |
|----------------|-----------|---------------------------------------------------------------------------------------|------------------------------------------------------------|--|--|--|
| CRM5342        | Synthetic | Μ                                                                                     | RNAi                                                       |  |  |  |
| Description    | siRNA     | to inhibit FOLH1 exp                                                                  | ression using RNA interference                             |  |  |  |
| Specificity    | FOLH      | 1 siRNA (Mouse) is a t                                                                | arget-specific 19-23 nt siRNA oligo duplexes designed to   |  |  |  |
|                | knock     | down gene expressio                                                                   | n.                                                         |  |  |  |
| Form           | Lyoph     | ilized powder                                                                         |                                                            |  |  |  |
| Gene Symbol    | FOLH      | FOLH1                                                                                 |                                                            |  |  |  |
| Alternative Na | ames MOPS | MOPSM; NAALAD1; Glutamate carboxypeptidase 2; Folate hydrolase 1;                     |                                                            |  |  |  |
|                | Folylp    | oly-gamma-glutamat                                                                    | e carboxypeptidase; FGCP; Glutamate carboxypeptidase       |  |  |  |
|                | II; GC    | PII; Membrane glutan                                                                  | nate carboxypeptidase; mGCP; N-acetylated-alpha-linked     |  |  |  |
|                | acidic    | dipeptidase I; NAALA                                                                  | Dase                                                       |  |  |  |
| Entrez Gene    | 53320     | 53320 (Mouse)                                                                         |                                                            |  |  |  |
| SwissProt      | 0354      | O35409 (Mouse)                                                                        |                                                            |  |  |  |
| Purity         | > 97%     | ,<br>)                                                                                |                                                            |  |  |  |
| Quality Contro | ol Oligoi | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure |                                                            |  |  |  |
|                | appro     | priate coupling efficie                                                               | ency. The oligo is subsequently purified by affinity-solid |  |  |  |
|                | phase     | extraction. The anne                                                                  | aled RNA duplex is further analyzed by mass                |  |  |  |
|                | spect     | rometry to verify the                                                                 | exact composition of the duplex. Each lot is compared to   |  |  |  |
|                | the pi    | revious lot by mass sp                                                                | ectrometry to ensure maximum lot-to-lot consistency.       |  |  |  |
| Components     | We of     | fers pre-designed set                                                                 | s of 3 different target-specific siRNA oligo duplexes of   |  |  |  |
|                | mous      | e FOLH1 gene. Each v                                                                  | ial contains 5 nmol of lyophilized siRNA. The duplexes can |  |  |  |
|                | be tra    | insfected individually                                                                | or pooled together to achieve knockdown of the target      |  |  |  |
|                | gene,     | which is most commo                                                                   | only assessed by qPCR or western blot.                     |  |  |  |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



## **Product Data Sheet**

| Component               | 15 nmol      | 30 nmol      |
|-------------------------|--------------|--------------|
| FOLH1 siRNA (Mouse) - A | 5 nmol x 1   | 5 nmol x 2   |
| FOLH1 siRNA (Mouse) - B | 5 nmol x 1   | 5 nmol x 2   |
| FOLH1 siRNA (Mouse) - C | 5 nmol x 1   | 5 nmol x 2   |
| Negative Control        | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water              | 1 ml x 1     | 1 ml x 2     |

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

| Plate   | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
|         | of medium    | of siRNA            |               | 2000         |
|         |              | 100 nM              | 0.5 μl        | 0.25 μl      |
| 96-well | 100 μl       | 50 nM               | 0.25 μl       | 0.25 μl      |
|         |              | 10 nM               | 0.05 μl       | 0.25 μl      |
|         |              | 100 nM              | 2.5 μl        | 1 µl         |
| 24-well | 500 μl       | 50 nM               | 1.25 μl       | 1 μΙ         |
|         |              | 10 nM               | 0.25 μl       | 1 μl         |
|         |              | 100 nM              | 5 μl          | 2 μΙ         |
| 12-well | 1 ml         | 50 nM               | 2.5 μl        | 2 μΙ         |
|         |              | 10 nM               | 0.5 μl        | 2 μΙ         |
| 6-well  | 2 ml         | 100 nM              | 10 µl         | 5 µl         |
|         |              | 50 nM               | 5 μl          | 5 μΙ         |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



## **Product Data Sheet**

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com