

Product Data Sheet

EIF3G siRNA (Mouse)

Source	Reactivity	Applications		
Synthetic	Μ	RNAi		
siRNA	to inhibit EIF3G expr	ession using RNA interference		
EIF3G	siRNA (Mouse) is a t	arget-specific 19-23 nt siRNA oligo duplexes designed to		
knock	down gene expression	on.		
Lyoph	ilized powder			
EIF3G	EIF3G			
nes EIF3P	EIF3P42; EIF3S4; Eukaryotic translation initiation factor 3 subunit G; eIF3g;			
Eukar	yotic translation initia	ation factor 3 RNA-binding subunit; eIF-3 RNA-binding		
subur	iit; Eukaryotic transla	tion initiation factor 3 subunit 4; eIF-3-delta; eIF3 p42;		
elF3 p	044			
53356	53356 (Mouse)			
Q9Z1I	D1 (Mouse)			
> 97%				
Oligor	nucleotide synthesis i	s monitored base by base through trityl analysis to ensure		
appro	priate coupling effici	ency. The oligo is subsequently purified by affinity-solid		
phase	extraction. The anne	ealed RNA duplex is further analyzed by mass		
specti	rometry to verify the	exact composition of the duplex. Each lot is compared to		
the pr	revious lot by mass sp	pectrometry to ensure maximum lot-to-lot consistency.		
We of	fers pre-designed set	s of 3 different target-specific siRNA oligo duplexes of		
mous	e EIF3G gene. Each vi	al contains 5 nmol of lyophilized siRNA. The duplexes can		
be tra	nsfected individually	or pooled together to achieve knockdown of the target		
gene,	which is most comm	only assessed by qPCR or western blot.		
	Synthetic siRNA EIF3G knock Lyoph EIF3G EIF3G Subur eIF3 p 53356 Q9Z11 > 97% Oligor appro phase spect the pr We of mous be tra	Synthetic M siRNA to inhibit EIF3G expr EIF3G siRNA (Mouse) is a tak knock down gene expression Lyophilized powder EIF3G EIF3P42; EIF3S4; Eukaryotic Eukaryotic translation initia subunit; Eukaryotic transla eIF3 p44 53356 (Mouse) Q9Z1D1 (Mouse) > 97% Oligonucleotide synthesis i appropriate coupling efficie phase extraction. The anne spectrometry to verify the the previous lot by mass sp We offers pre-designed set mouse EIF3G gene. Each vi be transfected individually		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

Component	15 nmol	30 nmol
EIF3G siRNA (Mouse) - A	5 nmol x 1	5 nmol x 2
EIF3G siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
EIF3G siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 µl	2 μΙ
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
6-well	2 ml	100 nM	10 µl	5 µl
o-well	2 111	50 nM	5 µl	5 μΙ

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com