

Product Data Sheet

VTI1A siRNA (Mouse)

Catalog #	Source	Reactivity	Applications			
CRM5385	Synthetic	Μ	RNAi			
Description	siRNA	to inhibit VTI1A expre	ession using RNA interference			
Specificity	VTI1A	VTI1A siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed to				
	knock	down gene expressio	n.			
Form	Lyoph	ilized powder				
Gene Symbol	VTI1A	VTI1A				
Alternative N	ames VTI1;	VTI1; VTI1L2; Vesicle transport through interaction with t-SNAREs homolog 1A;				
	Vesicl	e transport v-SNARE p	rotein Vti1-like 2; Vti1-rp2			
Entrez Gene	53611	L (Mouse)				
SwissProt	0891	O89116 (Mouse)				
Purity > 97%		,)				
Quality Contr	Control Oligonucleotide synthesis is monitored base by base through trityl analysis to e			sh trityl analysis to ensure		
	appro	priate coupling efficie	ncy. The oligo is subsequently p	urified by affinity-solid		
	phase	extraction. The anne	aled RNA duplex is further analy	zed by mass		
	spect	rometry to verify the e	exact composition of the duplex.	. Each lot is compared to		
	the pr	revious lot by mass sp	ectrometry to ensure maximum	lot-to-lot consistency.		
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of				
	mous	e VTI1A gene. Each via	al contains 5 nmol of lyophilized	siRNA. The duplexes can		
	be tra	insfected individually	or pooled together to achieve kr	nockdown of the target		
	gene,	gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent	15 nmol	30 nmol		
	VTI1	A siRNA (Mouse) - A	5 nmol x 1	5 nmol x 2		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

VTI1A siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
VTI1A siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
_		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
_		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 µl	5 μΙ
		10 nM	1 μΙ	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com