

Product Data Sheet

DNMT3L siRNA (Mouse)

Catalog #	Source	Reactivity	Applicat	ions	
CRM5524	Synthetic	Μ	RNAi		
Description	siRNA	to inhibit DNMT3L ex	pression using RNA inter	ference	
Specificity	DNM	DNMT3L siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed			
	to kno	ock down gene expres	ssion.		
Form	Lyoph	ilized powder			
Gene Symbol	DNM	DNMT3L			
Alternative N	ames DNA;	DNA; cytosine-5)-methyltransferase 3-like			
Entrez Gene	54427	7 (Mouse)			
SwissProt	Q9CW	/R8 (Mouse)			
Purity	> 97%	> 97%			
Quality Control Oligonucleotide synthesis is monitored base by base through trity		through trityl analysis to ensure			
	appro	priate coupling efficie	ency. The oligo is subsequ	ently purified by affinity-solid	
	phase	extraction. The anne	aled RNA duplex is furthe	r analyzed by mass	
	spect	rometry to verify the	exact composition of the	duplex. Each lot is compared to	
	the pr	revious lot by mass sp	ectrometry to ensure ma	ximum lot-to-lot consistency.	
Components	We of	fers pre-designed set	s of 3 different target-spe	cific siRNA oligo duplexes of	
	mous	e DNMT3L gene. Each	vial contains 5 nmol of ly	ophilized siRNA. The duplexes	
	can be	e transfected individu	ally or pooled together to	achieve knockdown of the	
target gene, which is most commonly assessed by qPCR or western b			PCR or western blot.		
	Com	ponent	15 nmol	30 nmol	
	DNM	1T3L siRNA (Mouse) -	A 5 nmol x 1	5 nmol x 2	
	DNM	1T3L siRNA (Mouse) -	B 5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

	DNMT3L siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
	Negative Control	2.5 nmol x 1	2.5 nmol x 2
_	DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com