

Product Data Sheet

SLC13A1 siRNA (Mouse)

Catalog #	Source	Reactivity	Applications		
CRM5596	Synthetic	Μ	RNAi		
Description	siRNA	to inhibit SLC13A1 e	pression using RNA interference		
Specificity	SLC13	SLC13A1 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed			
	to kno	ock down gene expre	ssion.		
Form	Lyoph	ilized powder			
Gene Symbol	SLC13	SLC13A1			
Alternative N	ames NAS1	NAS1; NASI1; Solute carrier family 13 member 1; NaSi-1; Renal sodium/sulfate			
	cotrai	nsporter; Na(+)/sulfat	e cotransporter		
Entrez Gene	55962	55961 (Mouse)			
SwissProt	Q9JHI	Q9JHI4 (Mouse)			
Purity	> 97%	, D			
Quality Control Oligonucleotide synthesis is monitored base by base through trityl a			sh trityl analysis to ensure		
	appro	priate coupling efficie	ency. The oligo is subsequently p	urified by affinity-solid	
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass			
	spect	rometry to verify the	exact composition of the duplex.	. Each lot is compared to	
	the p	revious lot by mass sp	ectrometry to ensure maximum	lot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	mous	e SLC13A1 gene. Each	vial contains 5 nmol of lyophiliz	ed siRNA. The duplexes	
	can b	e transfected individu	ally or pooled together to achiev	ve knockdown of the	
	target	target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	SLC1	.3A1 siRNA (Mouse) -	A 5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

SLC13A1 siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
SLC13A1 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
_		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
_		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
_		10 nM	0.5 μl	2 μΙ
		100 nM	10 µl	5 μΙ
6-well	2 ml	50 nM	5 µl	5 µl
		10 nM	1 μΙ	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com