

Product Data Sheet

TERF2IP siRNA (Mouse)

| Catalog # | Source | Reactivity | Applications | | |
|---------------|------------|---|---|-------------|--|
| CRM6003 | Synthetic | М | RNAi | | |
| Description | siRNA | to inhibit TERF2IP ex | pression using RNA interference | | |
| Specificity | TERF2 | 2IP siRNA (Mouse) is a | a target-specific 19-23 nt siRNA oligo duplexes | designed to | |
| | knock | down gene expression | on. | | |
| Form | Lyoph | nilized powder | | | |
| Gene Symbol | TERF2 | TERF2IP | | | |
| Alternative N | ames RAP1; | RAP1; Telomeric repeat-binding factor 2-interacting protein 1; TERF2-interacting | | | |
| | telom | eric protein 1; TRF2-i | interacting telomeric protein 1; Repressor/activ | vator | |
| | prote | in 1 homolog; RAP1 ዞ | nomolog | | |
| Entrez Gene | 57322 | 1 (Mouse) | | | |
| SwissProt | Q91V | Q91VL8 (Mouse) | | | |
| Purity | > 97% | 6 | | | |
| Quality Contr | ol Oligoi | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure | | | |
| | appro | opriate coupling effici | ency. The oligo is subsequently purified by affir | nity-solid | |
| | phase | e extraction. The anne | ealed RNA duplex is further analyzed by mass | | |
| | spect | rometry to verify the | exact composition of the duplex. Each lot is co | mpared to | |
| | the p | revious lot by mass sp | pectrometry to ensure maximum lot-to-lot con | sistency. | |
| Components | We of | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of | | | |
| | mous | e TERF2IP gene. Each | vial contains 5 nmol of lyophilized siRNA. The | duplexes | |
| | can b | e transfected individu | ually or pooled together to achieve knockdown | of the | |
| | target | target gene, which is most commonly assessed by qPCR or western blot. | | | |
| | Com | ponent | 15 nmol 30 nmol | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| TERF2IP siRNA (Mouse) - A | 5 nmol x 1 | 5 nmol x 2 |
|---------------------------|--------------|--------------|
| TERF2IP siRNA (Mouse) - B | 5 nmol x 1 | 5 nmol x 2 |
| TERF2IP siRNA (Mouse) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μl |
| | | 10 nM | 0.25 μl | 1 μl |
| | | 100 nM | 5 µl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| | | 100 nM | 10 µl | 5 μΙ |
| 6-well | 2 ml | 50 nM | 5 µl | 5 μl |
| | | 10 nM | 1 µl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



For research purposes only, not for human use

Product Data Sheet

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com