

# **Product Data Sheet**

## NUDT2 siRNA (Mouse)

Catalog #	Source	Reactivity	Applications		
CRM6613	Synthetic	Μ	RNAi		
Description	siRN	A to inhibit NUDT2 ex	pression using RNA interference		
Specificity	NUD	T2 siRNA (Mouse) is a	target-specific 19-23 nt siRNA oligo duplexes designed to		
	knoc	k down gene express	on.		
Form	Lyop	hilized powder			
Gene Symbol	NUD	NUDT2			
Alternative N	ames APAH	APAH1; Bis(5'-nucleosyl)-tetraphosphatase [asymmetrical]; Diadenosine			
	5'.5''	'-P1.P4-tetraphospha	te asymmetrical hydrolase; Ap4A hydrolase; Ap4Aase;		
	Diad	enosine tetraphospha	atase; Nucleoside diphosphate-linked moiety X motif 2;		
	Nudi	x motif 2			
Entrez Gene	6640	66401 (Mouse)			
SwissProt	P563	P56380 (Mouse)			
Purity	> 97	%			
Quality Contr	ol Oligo	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure			
	appr	opriate coupling effic	iency. The oligo is subsequently purified by affinity-solid		
	phas	e extraction. The ann	ealed RNA duplex is further analyzed by mass		
	spec	trometry to verify the	exact composition of the duplex. Each lot is compared to		
	the p	previous lot by mass s	pectrometry to ensure maximum lot-to-lot consistency.		
Components	We d	offers pre-designed se	ts of 3 different target-specific siRNA oligo duplexes of		
	mou	se NUDT2 gene. Each	vial contains 5 nmol of lyophilized siRNA. The duplexes		
	can l	be transfected individ	ually or pooled together to achieve knockdown of the		
	targe	et gene, which is most	commonly assessed by qPCR or western blot.		
A secold sectors from		Ale stand blat UI lasses	a historia de la la la successiva e flucence a successiva e de la successiva de la successiva de la successiva		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

Component	15 nmol	30 nmol
NUDT2 siRNA (Mouse) - A	5 nmol x 1	5 nmol x 2
NUDT2 siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
NUDT2 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 μΙ
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
6-well	2 ml	100 nM	10 µl	5 µl
		50 nM	5 μl	5 μΙ

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com