

Product Data Sheet

NDC80 siRNA (Mouse)

Catalog #	Source	Reactivity	Applications		
CRM6971	Synthetic	Μ	RNAi		
Description siRNA to inhibit NDC80 expression using RNA interference					
Specificity	NDC8	NDC80 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed to			
	knock	down gene expressi	on.		
Form	Lyoph	Lyophilized powder			
Gene Symbol	NDC8	NDC80			
Alternative N	ames HEC1;	HEC1; KNTC2; Kinetochore protein NDC80 homolog; Kinetochore protein Hec1;			
	Kineto	ochore-associated pro	otein 2		
Entrez Gene	67052	(Mouse)			
SwissProt	Q9D0	Q9D0F1 (Mouse)			
Purity >		> 97%			
Quality Control Oligonucleotide synthesis			is monitored base by base throug	h trityl analysis to ensure	
	appro	priate coupling effici	ency. The oligo is subsequently p	urified by affinity-solid	
	phase	extraction. The anne	ealed RNA duplex is further analy	zed by mass	
	spectr	ometry to verify the	exact composition of the duplex.	Each lot is compared to	
	the pr	evious lot by mass sp	pectrometry to ensure maximum	lot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	mouse	e NDC80 gene. Each	vial contains 5 nmol of lyophilized	d siRNA. The duplexes	
	can be	e transfected individu	ally or pooled together to achiev	e knockdown of the	
	target	target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	NDC	80 siRNA (Mouse) - A	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

NDC80 siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
NDC80 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 μl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
		100 nM	10 µl	5 μΙ
6-well	2 ml	50 nM	5 μl	5 μΙ
		10 nM	1 μl	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com