

Product Data Sheet

RNASEH2B siRNA (Mouse)

Catalog #	Source	Reactivity	Applications		
CRM7030	Synthetic	Μ	RNAi		
Description	siRNA	to inhibit RNASEH2	expression using RNA interfere	nce	
Specificity	RNAS	EH2B siRNA (Mouse)	is a target-specific 19-23 nt siRN	IA oligo duplexes designed	
	to kno	ock down gene expre	ssion.		
Form	Lyoph	ilized powder			
Gene Symbol	RNAS	RNASEH2B			
Alternative N	ames DLEU8	DLEU8; Ribonuclease H2 subunit B; RNase H2 subunit B; Deleted in lymphocytic			
	leukei	mia 8 homolog; Ribo	nuclease HI subunit B		
Entrez Gene	67153	3 (Mouse)			
SwissProt	Q80Z	Q80ZV0 (Mouse)			
Purity	> 97%	> 97%			
Quality Control Oligonucleotide synthesis is monitored base by base through tri			gh trityl analysis to ensure		
	appro	priate coupling effici	ency. The oligo is subsequently p	ourified by affinity-solid	
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass			
	specti	spectrometry to verify the exact composition of the duplex. Each lot is compared to			
	the pr	revious lot by mass s	pectrometry to ensure maximum	n lot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	mous	mouse RNASEH2B gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes			
	can be	can be transfected individually or pooled together to achieve knockdown of the			
	target	target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	RNA	SEH2B siRNA (Mouse) - A 5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

RNASEH2B siRNA (Mouse) - B	5 nmol x 1	5 nmol x 2
RNASEH2B siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 μl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
_		10 nM	0.5 μl	2 μΙ
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 μΙ
		10 nM	1 µl	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com