

Product Data Sheet

SEC14L2 siRNA (Mouse)

Catalog #	Source	Reactivity	Application	S	
CRM7360	Synthetic	М	RNAi		
Description	siRNA	A to inhibit SEC14L2 e	xpression using RNA interfere	ence	
Specificity	SEC14	SEC14L2 siRNA (Mouse) is a target-specific 19-23 nt siRNA oligo duplexes designed			
	to kn	ock down gene expre	ssion.		
Form	Lyopł	nilized powder			
Gene Symbol	SEC14	SEC14L2			
Alternative N	lames SEC14	SEC14-like protein 2; Alpha-tocopherol-associated protein; TAP			
Entrez Gene	6781	5 (Mouse)			
SwissProt	Q99J(08 (Mouse)			
Purity	> 97%	> 97%			
Quality Control Oligonucleotide synthesis is monitored base by base throug			rough trityl analysis to ensure		
	appro	opriate coupling effici	ency. The oligo is subsequent	ly purified by affinity-solid	
	phase	e extraction. The ann	ealed RNA duplex is further a	nalyzed by mass	
	spect	rometry to verify the	exact composition of the dup	plex. Each lot is compared to	
	the p	revious lot by mass s	pectrometry to ensure maxim	num lot-to-lot consistency.	
Components	We o	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	mous	e SEC14L2 gene. Eacl	n vial contains 5 nmol of lyopl	hilized siRNA. The duplexes	
	can b	e transfected individ	ually or pooled together to ac	hieve knockdown of the	
	target gene, which is most commonly assessed by qPCR or western blot.			or western blot.	
	Com	ponent	15 nmol	30 nmol	
	SEC	14L2 siRNA (Mouse) -	A 5 nmol x 1	5 nmol x 2	
	SEC	14L2 siRNA (Mouse) -	B 5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

SEC14L2 siRNA (Mouse) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com